

Small x behavior of parton distributions.

Analytical and “frozen” coupling constants.

BFKL corrections

OUTLINE

1. Introduction
2. Results: flat initial conditions for PDFs at small x values
(i.e. *min* information about initial conditions, or
min contribution from initial conditions.)
3. Conclusions and Prospects

1. Introduction to DIS

A. Deep-inelastic scattering cross-section:

$$\sigma \sim L^{\mu\nu} F^{\mu\nu}$$

Hadron part $F^{\mu\nu}$ ($Q^2 = -q^2 > 0$, $x = Q^2/[2(pq)]$):

$$F^{\mu\nu} = \left(-g^{\mu\nu} + \frac{q^\mu q^\nu}{q^2}\right) F_1(x, Q^2) \\ - \left(p^\mu - \frac{(pq)}{q^2} q^\mu\right) \left(p^\nu - \frac{(pq)}{q^2} q^\nu\right) \frac{2x}{q^2} F_2(x, Q^2) + \dots,$$

where $F_k(x, Q^2)$ ($k = 1, 2, 3, L$) - are DIS SF and q and p are photon and hadron (parton) momentums.

B. Wilson operator expansion: Mellin moments $M_k(n, Q^2)$ of DIS SF $F_k(x, Q^2)$ can be represented as sum

$$M_k(n, Q^2) = \sum_{a=NS, SI, g} \underbrace{C_k^a(n, Q^2/\mu^2)}_{\text{Coeff. function}} A_a(n, \mu^2),$$

where $A_a(n, \mu^2) = \langle N | \mathcal{O}_{\mu_1, \dots, \mu_n}^a | N \rangle$ are matrix elements of the Wilson operators $\mathcal{O}_{\mu_1, \dots, \mu_n}^a$.

C. The matrix elements $A_a(n, \mu^2)$ are Mellin moments of the unpolarized PD $f_a(n, \mu^2)$.

DGLAP [= Renormgroup] equations:

$$\frac{d}{d \ln Q^2} f_a(x, Q^2) = \int_x^1 \frac{dy}{y} \sum_b W_{b \rightarrow a}(x/y) f_b(y, Q^2). \quad (1)$$

The anomalous dimensions (AD) $\gamma_{ab}(n)$ of the twist-2 Wilson operators $\mathcal{O}_{\mu_1, \dots, \mu_n}^a$ (hereafter $a_s = \alpha_s/(4\pi)$)

$$\gamma_{ab}(n) = \int_0^1 dx x^{n-1} W_{b \rightarrow a}(x) = \sum_{m=0}^{\infty} \gamma_{ab}^{(m)}(n) a_s^m,$$

All parton densities are multiplied by x , t.e.

structure function = combination of parton densities.

3. Method

(C.Lopez and F.J.Yndurain, 1980,1981), (A.V.K., 1994)

Here I present briefly the method, which leads to the possibility to replace the Mellin convolution of two functions

$$f_1(x) \otimes f_2(x) \equiv \int_x^1 \frac{dy}{y} f_1(y) f_2(x/y)$$

by a simple products at small x .

A. So, if $f_1(x) = B_k(x, Q^2)$ is perturbatively calculated Wilson kernel and $f_2(x) = x f_a(x, Q^2) \sim x^{-\delta}$ at $x \rightarrow 0$, then

$$f_1(x) \otimes f_2(x) \approx M_k(1 + \delta, Q^2) f_2(x) \quad (2)$$

where $M_k(1 + \delta, Q^2)$ is the analytical continuation to non-integer arguments of the Mellin moment $M_k(n, Q^2)$ of $B_k(x, Q^2)$:

$$M_k(n, Q^2) = \int_0^1 x^{n-2} B_k(x, Q^2) \quad (3)$$

The equation (2) is correct if the moment $M_k(n, Q^2)$ has no singularity at $n \rightarrow 1$.

B. The general case

($M(n)$ contains the singularity at $n \rightarrow 1$):

the form of subasymptotics of $f_2(x)$ starts to be important.

Let PD have the different forms:

- Regge-like form $xf_R(x) = x^{-\delta}\tilde{f}(x)$,
- Logarithmic-like form $xf_L(x) = x^{-\delta}\ln(1/x)\tilde{f}(x)$,
- Bessel-like form $xf_I(x) = x^{-\delta}I_k(2\sqrt{\hat{d}\ln(1/x)})\tilde{f}(x)$,

where $\tilde{f}(x)$ and its derivative $\tilde{f}'(x) \equiv d\tilde{f}(x)/dx$ are smooth at $x = 0$ and both are equal to zero at $x = 1$:

$$\tilde{f}(1) = \tilde{f}'(1) = 0$$

Then ($i = R, L, I$)

$$f_1(x) \otimes f_2(x) \approx \tilde{M}_k(1 + \delta_i, Q^2) f_2(x),$$

where $\tilde{M}_{1+\delta_i} = M_{1+\delta}$ with $1/\delta \rightarrow 1/\tilde{\delta}_i$.

Regge-like behavior:

$$1/\tilde{\delta}_R = 1/\delta \left[1 - x^\delta \frac{\Gamma(1 - \delta)\Gamma(\nu)}{\Gamma(1 + \nu - \delta)} \right],$$

where $x f_R(x) \sim (1 - x)^\nu$ at $x \rightarrow 1$.

The second term comes from low part of convolution integral

$$f_1(x) \otimes f_2(x) \equiv \int_x^1 \frac{dy}{y} f_1(x/y) f_2(y) \quad (4)$$

So,

$$\frac{1}{\tilde{\delta}_R} = \frac{1}{\delta} \quad \text{if} \quad x^\delta \ll 1$$

and

$$\frac{1}{\tilde{\delta}_R} = \ln \frac{1}{x} - [\Psi(1 + \nu) - \Psi(1)] \quad \text{if} \quad \delta = 0$$

Analogously, for nonRegge behavior at $\delta \rightarrow 0$

$$\frac{1}{\tilde{\delta}_L} = \frac{1}{2} \ln \frac{1}{x} + O(1/\ln(1/x)),$$
$$\frac{1}{\tilde{\delta}_I} = \sqrt{\frac{\ln(1/x)}{\hat{d}}} \frac{I_{k+1}(2\sqrt{\hat{d}\ln(1/x)})}{I_k(2\sqrt{\hat{d}\ln(1/x)})}$$

3. Generalized double-logarithmic approach

(A.V.K. and G.Parente, 1998),

(A.Yu.Illarionov, A.V.K. and G.Parente, 2004)

(Generalized double asymptotic scaling)

1 Leading order without quarks (a pedagogical example)

At the momentum space, the solution of the DGLAP equation is

$$M_g(n, Q^2) = M_g(n, Q_0^2) e^{-d_{gg}(n)s},$$

where $M_g(n, Q^2)$ are the moments of the gluon distribution,

$$s = \ln \left(\frac{a_s(Q_0^2)}{a_s(Q^2)} \right), \quad a_s(Q^2) = \frac{\alpha_s(Q^2)}{4\pi} \quad \text{and} \quad d_{gg} = \frac{\gamma_{gg}^{(0)}(n)}{2\beta_0}$$

The terms $\gamma_{gg}^{(0)}(n)$ and β_0 are respectively the LO coefficients of the gluon-gluon AD and the QCD β -function.

For any perturbatively calculable variable $Q(n)$, it is very convenient to separate the singular part when $n \rightarrow 1$ (denoted by “ \widehat{Q} ”) and the regular part (marked as “ \overline{Q} ”):

$$Q(n) = \frac{\widehat{Q}}{n-1} + \overline{Q}(n)$$

Then, the above equation can be represented by the form

$$M_g(n, Q^2) = M_g(n, Q_0^2) e^{-\hat{d}_{gg} s_{LO}/(n-1)} e^{-\bar{d}_{gg}(n) s_{LO}},$$

with $\hat{\gamma}_{gg} = -8C_A$ and $C_A = N$ for $SU(N)$ group.

Finally, if one takes the flat boundary conditions (i.e. *min* information about initial conditions, or *min* contribution from initial conditions.)

$$x f_a(x, Q_0^2) = A_a, \quad \rightarrow \quad M_a(n, Q_0^2) = \frac{A_a}{n-1} \quad (5)$$

1.1 Classical double-logarithmic case ($\bar{d}_{gg}(n) = 0$)

(A.D.Rujula, S.L.Glashow, H.D.Politzer, S.B.Treiman, F.Wilczek and A.Zee, 1974)

Then, expanding the second exponential in the above equation

$$M_g^{cdl}(n, Q^2) = A_g \sum_{k=0}^{\infty} \frac{1}{k!} \frac{(-\hat{d}_{gg} s_{LO})^k}{(n-1)^{k+1}}$$

and using the Mellin transformation for $(\ln(1/x))^k$:

$$\int_0^1 dx x^{n-2} (\ln(1/x))^k = \frac{k!}{(n-1)^{k+1}}$$

we immediately obtain the well known double-logarithmic behavior

$$f_g^{cdl}(x, Q^2) = A_g \sum_{k=0}^{\infty} \frac{1}{(k!)^2} (-\hat{d}_{gg} s_{LO})^k (\ln(1/x))^k = A_g I_0(\sigma_{LO}),$$

where $I_0(\sigma_{LO})$ is the modified Bessel function with argument $\sigma_{LO} = 2\sqrt{\hat{d}_{gg} s_{LO} \ln(x)}$. (R.D.Ball and S.Forte, 1994),

1.2 The more general case

For a regular kernel $\tilde{K}(x)$, having Mellin moment
(nonsingular at $n \rightarrow 1$)

$$K(n) = \int_0^1 dx x^{n-2} \tilde{K}(x)$$

and the PD $f_a(x)$ in the form $I_\nu(\sqrt{\hat{d} \ln(1/x)})$ we have the following equation

$$\tilde{K}(x) \otimes f_a(x) = K(1) f_a(x) + O\left(\sqrt{\frac{\hat{d}}{\ln(1/x)}}\right)$$

So, one can find the general solution for the LO gluon density without the influence of quarks

$$f_g(x, Q^2) = A_g I_0(\sigma_{LO}) e^{-\bar{d}_{gg}(1) s_{LO}} + O(\rho_{LO}),$$

where (R.D.Ball and S.Forte, 1994)

$$\rho_{LO} = \sqrt{\frac{\hat{d}_{gg} s_{LO}}{\ln(x)}} = \frac{\sigma_{LO}}{2 \ln(1/x)}, \quad \bar{\gamma}_{gg}^{(0)}(1) = 22 + \frac{4}{3}f$$

and

$$\bar{d}_{gg}(1) = 1 + \frac{4f}{3\beta_0}$$

with f as the number of active quarks.

2 Leading order (complete)

At the momentum space, the solution of the DGLAP equation at LO has the form (*after diagonalization*)

$$M_a(n, Q^2) = M_a^+(n, Q^2) + M_a^-(n, Q^2) \quad \text{and}$$

$$M_a^\pm(n, Q^2) = M_a^\pm(n, Q_0^2) e^{-d_\pm(n)s} = M_a^\pm e^{-\hat{d}_\pm s / (n-1)} e^{-\bar{d}_\pm(n)s},$$

where

$$M_a^\pm(n, Q^2) = \varepsilon_{ab}^\pm(n) M_b(n, Q^2), \quad d_{ab} = \frac{\gamma_{ab}^{(0)}(n)}{2\beta_0},$$

$$d_\pm(n) = \frac{1}{2}[(d_{gg}(n) + d_{qq}(n))$$

$$\pm (d_{gg}(n) - d_{qq}(n)) \sqrt{1 + \frac{4d_{qg}(n)d_{gq}(n)}{(d_{gg}(n) - d_{qq}(n))^2}}]$$

$$\varepsilon_{qq}^\pm(n) = \varepsilon_{gg}^\mp(n) = \frac{1}{2} \left(1 + \frac{d_{qq}(n) - d_{gg}(n)}{d_\pm(n) - d_\mp(n)} \right),$$

$$\varepsilon_{ab}^{\pm}(n) = \frac{d_{ab}(n)}{d_{\pm}(n) - d_{\mp}(n)} (a \neq b)$$

As the singular (when $n \rightarrow 1$) part of the + component of the anomalous dimension is !!! $\hat{d}_+ = \hat{d}_{gg} = -4C_A/\beta_0$!!! while the - component does not exist: !!! $(\hat{d}_- = 0)$!!! , we consider below both cases separately.

2.1 The “+” component

The analysis of the “+” component is practically identical to the case studied before. The only difference lies in the appearance of new terms $\varepsilon_{ab}^+(n)$!!! . If they are expanded in the vicinity of $n = 1$ in the form $\varepsilon_{ab}^+(n) = \bar{\varepsilon}_{ab}^+ + (n - 1)\tilde{\varepsilon}_{ab}^+$, !!! then for the terms $\bar{\varepsilon}_{ab}^+$ multiplying $M_b(n, Q^2)$, we have the same results as in previous section:

$$\bar{\varepsilon}_{ab}^+ M_b(n, Q^2) \xrightarrow{\mathcal{M}^{-1}} \bar{\varepsilon}_{ab}^+ A_b I_0(\sigma_{LO}) e^{-\bar{d}_+(1)s_{LO}} + O(\rho_{LO}),$$

where the symbol $\xrightarrow{\mathcal{M}^{-1}}$ denotes the inverse Mellin transformation.

The values of σ and ρ coincide with those defined in the previous section because $\hat{d}_+ = \hat{d}_{gg}$.

The terms $\tilde{\varepsilon}_{ab}^+$ that come with the additional factor $(n - 1)$ in front, lead to the following results

$$(n - 1)\tilde{\varepsilon}_{ab}^+ \frac{A_b}{(n - 1)} e^{-\hat{d}_+ s_{LO}/(n-1)} = \tilde{\varepsilon}_{ab}^+ A_b \sum_{k=0}^{\infty} \frac{1}{k!} \frac{(-\hat{d}_+ s_{LO})^k}{(n - 1)^k}$$

$$\xrightarrow{\mathcal{M}^{-1}} \tilde{\varepsilon}_{ab}^+ A_b \sum_{k=0}^{\infty} \frac{1}{k!} \frac{1}{(k - 1)!} (-\hat{d}_+ s_{LO})^k (\ln(1/x))^{k-1}$$

$$= \tilde{\varepsilon}_{ab}^+ A_b \rho_{LO} I_1(\sigma_{LO}),$$

i.e. the additional factor $(n - 1)$ in momentum space leads to replacing the Bessel function $I_0(\sigma_{LO})$ by $\rho_{LO} I_1(\sigma_{LO})$ in x -space.

Thus, we obtain that the term $\varepsilon_{ab}^+(n) M_b(n, Q^2)$ leads to the following contribution in x space **!!!** :

$$(\bar{\varepsilon}_{ab}^+ I_0(\sigma_{LO}) + \tilde{\varepsilon}_{ab}^+ \rho_{LO} I_1(\sigma_{LO})) A_b e^{-\bar{d}_+(1) s_{LO}} + O(\rho_{LO})$$

Because the Bessel function $I_\nu(\sigma)$ has the ν -independent asymptotic behavior **!!!** $e^\sigma/\sqrt{\sigma}$ at $\sigma \rightarrow \infty$ (i.e. $x \rightarrow 0$), the second term is $O(\rho)$ and must be kept only **!!!** when $\bar{\varepsilon}_{ab}^+ = 0$. This is the case for the quark distribution at the LO approximation.

Using the concrete AD values, one has

$$f_g^+(x, Q^2) = (A_g + \frac{4}{9}A_q)I_0(\sigma_{LO})e^{-\bar{d}_+(1)s_{LO}} + O(\rho_{LO}) \quad \text{and}$$

$$f_q^+(x, Q^2) = \frac{f}{9}(A_g + \frac{4}{9}A_q)\rho_{LO}I_1(\sigma_{LO})e^{-\bar{d}_+(1)s_{LO}} + O(\rho_{LO})$$

where $\bar{d}_+(1) = 1 + 20f/(27\beta_0)$.

2.2 the “-” component

In this case the anomalous dimension is regular !!! and one has

$$\varepsilon_{ab}^-(n) A_b e^{-d_-(n)s} \xrightarrow{\mathcal{M}^{-1}} \bar{\varepsilon}_{ab}^-(1) A_b e^{-d_-(1)s_{LO}} + O(x)$$

Using the concrete AD values !!! , we have

$$f_g^-(x, Q^2) = -\frac{4}{9} A_q e^{-d_-(1)s_{LO}} + O(x) \text{ and}$$

$$f_q^-(x, Q^2) = A_q e^{-d_-(1)s_{LO}} + O(x),$$

where $d_-(1) = 16f/(27\beta_0)$.

Finally we present the full small x asymptotic results for PD and F_2 structure function at LO of perturbation theory:

$$f_a(x, Q^2) = f_a^+(x, Q^2) + f_a^-(x, Q^2) \quad \text{and}$$
$$F_2(x, Q^2) = e \cdot f_q(z, Q^2)$$

where f_q^+, f_g^+, f_q^- and f_g^- were already given before and $e = \frac{\sum_1^f e_i^2}{f}$ is the average charge square of the f active quarks.

Extension to NLO is trivial and can be found in (A.V.K. and G.Parente, 1998)

So, we resume the steps we have followed to reach the small x approximate solution of DGLAP shown above:

- Use the n -space exact solution.
- Expand the perturbatively calculated parts (AD and coefficient functions) in the vicinity of the point $n = 1$.
- The singular part with the form

$$A_a(n - 1)^k e^{-\hat{d}s_{LO}/(n-1)}$$

leads to Bessel functions in the x -space in the form

$$A_a \left(\frac{\hat{d}s_{LO}}{\ln x} \right)^{(k+1)/2} I_{k+1} \left(2 \sqrt{\hat{d}s_{LO} \ln x} \right)$$

- The regular part $B(n) \exp(-\bar{d}(n)s_{LO})$ leads to the additional coefficient

$$B(1)\exp(-\bar{d}(1)s_{LO}) + O(\sqrt{\hat{d}s_{LO}/\ln x})$$

behind of the Bessel function **!!!** in the x -space. Because the accuracy is $O(\sqrt{\hat{d}s_{LO}/\ln x})$, it is necessary to use only the first nonzero term **!!!**, i.e. all terms $(n-1)^k$ in front of $\exp(-\hat{d}/(n-1))$, with the exception of one with the smaller k value, can be neglected.

- If the singular part at $n \rightarrow 1$ is absent, i.e. $\hat{d} = 0$, the result in the x -space is determined by $B(1)\exp(-\bar{d}(1)s_{LO})$ with accuracy $O(x)$.

4. Fits of HERA data

At low x , the structure function $F_2(x, Q^2)$ is related to parton densities as (A.V.K. and G.Parente, 1998)

at LO

$$F_2(x, Q^2) = \frac{5}{18} f_q(x, Q^2)$$


at NLO


$$F_2(x, Q^2) = \frac{5}{18} \left[f_q(x, Q^2) + \frac{2f}{3} a_s(Q^2) f_g(x, Q^2) \right].$$


Fits of HERA experimental data of the structure function $F_2(x, Q^2)$ (A.Yu.Illarionov, A.V.K. and G.Parente, 2004)

!!! Only three parameters: Q_0^2 , A_q and A_g

Λ_{QCD} cannot be extracted in small x Physics.







5. Analytical and “frozen” coupling constants

Two modifications of the coupling constant (G.Cvetic, A.Yu.Illarinov, B.A. Kniehl, and A.V.Kotikov, 2009)

A. More phenomenological.

(G.Curci, M.Greco and Y.Sristava, 1979), (M.Greco, G. Penso and Y.Sristava, 1980), (N.N.Nikolaev and B.M.Zakharov, 1991,1992), (B.Badelek, J.Kwiecinski and A.Stasto, 1997), (A.M.Badalian and Yu.A.Simonov, 1997)

We introduce freezing of the coupling constant by changing its argument $Q^2 \rightarrow Q^2 + M_\rho^2$, where M_ρ is usually the ρ -meson mass. Thus, in the formulae of the previous Sections we should do the following replacement

$$a_s(Q^2) \rightarrow a_{fr}(Q^2) \equiv a_s(Q^2 + M_\rho^2) \quad (6)$$

B. Theoretical approach.

Incorporates the Shirkov-Solovtsov idea (D.V.Shirkov and L.I.Solovtsov, 1997), about analyticity of the coupling constant that leads to the additional its power dependence.

(K.A.Milton, A.V. Nesterenko, O.Solovtsova, G. Cvetič,
C. Valenzuela, I. Schmidt, O. Teryaev, N. Stefanis, A. Bakulev,
S. Mikhailov, ...)

Then, in the formulae of the previous Section the coupling constant $a_s(Q^2)$ should be replaced as follows

$$a_{an}^{LO}(Q^2) = a_s(Q^2) - \frac{1}{\beta_0} \frac{\Lambda_{LO}^2}{Q^2 - \Lambda_{LO}^2} \quad (7)$$

at the LO approximation and

$$a_{an}(Q^2) = a_s(Q^2) - \frac{1}{2\beta_0} \frac{\Lambda^2}{Q^2 - \Lambda^2} + \dots \quad (8)$$

at the NLO approximation, where the symbol ... marks numerically small terms.




Table 1: The result of the LO and NLO fits to H1 and ZEUS data for different low Q^2 cuts. In the fits f is fixed to 4 flavors.

	A_g	A_q	Q_0^2 [GeV 2]	$\chi^2/n.o.p.$
$Q^2 \geq 1.5\text{GeV}^2$				
LO	0.784 ± 0.016	0.801 ± 0.019	0.304 ± 0.003	754/609
LO&an.	0.932 ± 0.017	0.707 ± 0.020	0.339 ± 0.003	632/609
LO&fr.	1.022 ± 0.018	0.650 ± 0.020	0.356 ± 0.003	547/609
NLO	-0.200 ± 0.011	0.903 ± 0.021	0.495 ± 0.006	798/609
NLO&an.	0.310 ± 0.013	0.640 ± 0.022	0.702 ± 0.008	655/609
NLO&fr.	0.180 ± 0.012	0.780 ± 0.022	0.661 ± 0.007	669/609
$Q^2 \geq 0.5\text{GeV}^2$				
LO	0.641 ± 0.010	0.937 ± 0.012	0.295 ± 0.003	1090/662
LO&an.	0.846 ± 0.010	0.771 ± 0.013	0.328 ± 0.003	803/662
LO&fr.	1.127 ± 0.011	0.534 ± 0.015	0.358 ± 0.003	679/662
NLO	-0.192 ± 0.006	1.087 ± 0.012	0.478 ± 0.006	1229/662
NLO&an.	0.281 ± 0.008	0.634 ± 0.016	0.680 ± 0.007	633/662
NLO&fr.	0.205 ± 0.007	0.650 ± 0.016	0.589 ± 0.006	670/662


- Usage of the analytical and “frozen” coupling constants leads to improvement with data: χ^2 decreased twicely
- Really, no difference between results based on the analytical and “frozen” coupling constants.


!!! One example of application the analytical and “frozen” coupling constants: (A.V.Kotikov, A.V.Lipatov and N.P.Zotov, 2004)


New *H1&ZEUS* (2010) experimental data for F_2 : there is a good agreement for $Q^2 \geq 2 \text{ GeV}^2$.
Fits with $Q^2 \geq 0.5 \text{ GeV}^2$ are in progress.

The results for F_2 and for the slope of the SF F_2
The double-logarithmic behaviour can mimic a power law shape
over a limited region of x, Q^2 .

$$f_a(x, Q^2) \sim x^{-\lambda_a^{eff}(x, Q^2)} \quad \text{and} \quad F_2(x, Q^2) \sim x^{-\lambda_{F_2}^{eff}(x, Q^2)}$$







6. BFKL corrections

Idea:

(in this section it is convenient to replacy $\gamma^{\text{DIS}}/2 \rightarrow -\gamma$)

$$\gamma_+(n) = \frac{\hat{\gamma}_+}{\omega} + \bar{\gamma}_+(n), \quad (\omega = n - 1)$$

$$\frac{\hat{\gamma}_+}{\omega} \rightarrow \gamma_{\text{BFKL}}(\omega) = \gamma(\omega)$$

$$\omega = 4A_s \chi(\gamma), \quad \chi(\gamma) = 2\Psi(1) - \Psi(\gamma) - \Psi(1 - \gamma),$$

$$A_s = C_A a_s, \quad \Psi(\gamma) = \frac{d(\ln(\Gamma(\gamma)))}{d\gamma}$$

I. *Expansions*

$$\chi(\gamma) = \frac{1}{\gamma} + 2 \sum_{k=1}^{\infty} \zeta(2k+3) \gamma^{2k+2},$$

that is equal

$$\gamma = \gamma_0 \left[1 + 2 \sum_{k=1}^{\infty} \zeta(2k+3) \gamma_0^{2k+3} \right], \quad \gamma_0 = \frac{4A_s}{\omega} \quad (9)$$

or equal to

$$\gamma = \gamma_0 + \sum_{k=3}^{\infty} C_k \gamma_0^{k+1}$$

with the following nonzero C_k :

$$C_3 = 2\zeta(3), C_5 = 2\zeta(5), C_6 = 12\zeta^2(3), C_7 = 2\zeta(7), \dots \quad (10)$$

For the renormalization exponent we have the following replacement

$$\begin{aligned} & \exp\left\{-\frac{1}{\beta_0} \int_{a_s(Q_0^2)}^{a_s(Q^2)} \frac{da}{a^2} \frac{\hat{\gamma}_+(a)}{\omega}\right\} \\ \rightarrow & \exp\left\{-\frac{1}{\beta_0} \int_{a_s(Q_0^2)}^{a_s(Q^2)} \frac{da}{a^2} \gamma(a)\right\} \end{aligned}$$

and the following additional contribution

$$-\frac{1}{\beta_0} \int \frac{da}{a^2} \sum_{k=3}^{\infty} C_k \gamma_0^{k+1}(a) = -\frac{1}{\beta_0} \frac{1}{a} \sum_{k=3}^{\infty} \frac{C_k}{k} \gamma_0^{k+1}(a)$$

So, there is the additional factor k in denominator. We hope that in this form the BFKL corrections will be not so large!!!

II. *Exact form*

So, we should consider the contribution

$$\exp\left\{-\frac{1}{\beta_0} \int_{a_s(Q_0^2)}^{a_s(Q^2)} \frac{da}{a^2} \gamma(a)\right\}$$

Using integration by parts procedure:

$$\int \frac{da}{a^2} \gamma(a) = -\frac{1}{a} \gamma(a) + \int \frac{da}{a} \frac{d\gamma(a)}{da} = -\frac{1}{a} \gamma(a) + \int \frac{d\gamma}{a}$$

BFKL:

$$\frac{1}{a} = \frac{4C_A}{\omega} \chi(\gamma),$$

and

$$\int d\gamma \chi(\gamma) = 2\Psi(1)\gamma + \ln \frac{\Gamma(1-\gamma)}{\Gamma(\gamma)}$$

So, we have

$$\exp \left\{ -\frac{1}{\beta_0} \int_{a_s(Q_0^2)}^{a_s(Q^2)} \frac{da}{a^2} \gamma(a) \right\} = \frac{R(Q^2)}{R(Q_0^2)},$$

where

$$R(Q^2) = \left[\frac{\Gamma(\gamma)}{\Gamma(1-\gamma)} \right]^{d_0} \exp \{ -d_0 \gamma (\Psi(\gamma) + \Psi(1-\gamma)) \} \quad (11)$$

with

$$d_0 = \frac{4C_A}{\beta_0 \omega} = -\frac{\hat{d}_+}{\omega} = -\frac{\hat{d}_{gg}}{\omega}$$

At $a_s \rightarrow 0$ (i.e. $\gamma \rightarrow 0$) we recover the singular part of the LO contributions

$$R(Q^2) \rightarrow \gamma^{-d_0} \rightarrow a_s^{-d_0}$$

The following steps:

- To perform the inverse Mellin transformation
(in some approximate form,
because the structure of (11) is rather complicated).
- To perform fits with the terms (11) taken into account.

Conclusion

- I have demonstrated the low x asymptotics of parton densities and SF F_2 .
- Low x asymptotics of F_2 are in good agreement with data from HERA at $Q^2 \geq 2.5 \text{ GeV}^2$.
- Usage of the analytical and “frozen” coupling constants leads to improvement with data from HERA at $Q^2 \leq 2.5 \text{ GeV}^2$.
- BFKL corrections taken into account (only first steps).

Next steps:

- To add the NNLO corrections (which has $\sim 1/(n-1)^2$ poles at $n \rightarrow 1$). So, the NNLO small- x asymptotics $\sim \exp[\sim (\ln(1/x))^{2/3}]$ is more singular than the corresponding LO and NLO ones $\sim \exp[\sim \sqrt{\ln(1/x)}]$.
- To finish with BFKL corrections (at LO and NLO levels of approximation).