

QCD results from DZero

Dmitry Bandurin

XXII Baldin Seminar on HEP Problems
September 19, Dubna, Russia

Outline

- ◆ Jet production:
 - Inclusive jets
 - Dijets
 - 3-jets
- ◆ $V + \text{jets}$ production
 - $V + \text{inclusive jets}$
 - $V + \text{heavy flavor jets}$
- ◆ Inclusive photon and di-photon production
- ◆ Underlying events and double parton interactions

Selected some of the latest results. More results can be found on:


<http://www-d0.fnal.gov/Run2Physics/WWW/results/qcd.htm>


Also, a detailed overview of QCD measurements at the Tevatron:

1409.xxxx [hep-ex]


Should be published in Rev.Mod.Phys. A soon.

Tools








Jet Results



PDF, α_s
searches for New Physics




Motivations for jet measurements

- providing constraint on PDF:
 - x - Q^2 regions accessible at fixed target, DIS, Tevatron and LHC are complementary to each other
- α_s extraction
- studying internal jet substructure
- searches for new phenomena are limited without proper understanding QCD background



Jet Energy Scale

- Data and theory are corrected to the particle level: very challenging experimental issue, especially JES
- Getting precise (1-2%) JES results takes time. See detailed description at NIM A763, 442 (2014).




Inclusive jet production


- Inclusive is one of the most elementary measurements at hadron colliders.
- Inclusive jet cross sections at Tevatron test pQCD over 8-9 orders of magnitude up to 0.7 TeV
- **Primary and powerful source of PDF constraint!**



DØ, Cone





CDF, kT



Constraints on PDF


D0&CDF jet data favored lower bound of the theoretical (CTEQ6.5M PDF) predictions, with smaller gluon content at high x . Experimental uncertainties at high p_T are lower than theoretical (largely PDF ones):
 => constrain PDF

PRD75, 092006 (2007), PRL 101, 062001 (2008), PRD85, 052006 (2012)




The measurements lead to modified gluon PDF (especially at $x > 0.25$) and reduced PDF uncertainties.

MSTW 2008 uses CDF kT and D0 cone results.



Dijet mass cross section measurement




- Measurement of dijet mass in six rapidity bins, $|y|_{\text{max}} = \max(|y_1|, |y_2|)$

Non-perturbative corrections (-10%, 23%)

Comparison to NLO pQCD with MSTW2008 and CTEQ6.6M NLO PDFs,

$$\mu_F = \mu_R = (p_{T1} + p_{T2})/2$$






Last mass bin is at $\sim 1.3 \text{ TeV}$!

- 40—60% difference between PDFs (MSTW2008/CTEQ6.6) at high masses
- Data/QCD in good agreement in central region
- Data are lower than central pQCD prediction at higher rapidities

Three jet mass cross section

- Good agreement seen between data and NLO (MSTW2008) for all cases.
- Comparisons to ABKM09, NNPDF2.1, HERA1.0 are also provided.
- χ^2 test is done for 3 theor. scales and all α_s values available for a given PDF set
- *Best χ^2 results for MSTW2008, NNPDF2.1*


χ^2 test for *central* PDF values



$R_{3/2}$ results

Ratio of inclusive 3 to 2 jet cross sections


Phys. Lett. B 720, 6 (2013)



- Good agreement everywhere
 - Some shape of data/theory at lowest $p_{Tmin} = 30$ GeV
 - Best agreement is at scales= p_{Tmax} for $p_{Tmin} \geq 50$ GeV
 - The results for CT10 and NNPDFv2.1 PDFs agree with those obtained for MSTW2008NLO to better than 0.4%
- => open road for α_s extraction**

Azimuthal decorrelations


Phys. Lett. B 721, 212 (2013)



- $R_{\Delta\phi}(H_T, y^*, \Delta\phi_{\max})$
- First measurement of rapidity dependence of Dijet azimuthal decorrelations



$1/k = \text{LO/NLO ratio}$





- Good agreement where non-pert. Corrections, and LO/NLO ratio are not too large

Strong Coupling Constant

α_s is the fundamental QCD quantity

α_s is the least known of the couplings ($\Delta\alpha_s$ (WA) = 0.6%)

α_s has influence on GUT and $\Delta\alpha_s$ translates into uncertainty on PDFs and cross sections



- Renormalization Group Equation (RGE) relates α_s values at different scales (Q)
- Predictions tested at LEP, HERA up to $Q \approx 208$ GeV, +recently by Tevatron from inclusive Jets (54-145 GeV)



Strong Coupling Constant

α_s is the fundamental QCD quantity

α_s is the least known of the couplings ($\Delta\alpha_s$ (WA) = 0.6%)

α_s has influence on GUT and $\Delta\alpha_s$ translates into uncertainty on PDFs and cross sections

=> Jet data can be used to extract the running of α_s



Running of $\alpha_s(Q)$ can be modified for large Q , e.g. by extra dimensions
=> should be tested using variable free of RGE (PDF) dependence




Strong Coupling Constant

Phys. Lett. B 721, 212(2013)

New variable introduced in D0 that characterizes angular correlations of jets and gives an average number of neighboring jets around a reference jet, measured triple differentially: $R_{\Delta R}(p_T, \Delta R, p_{T\text{-nbr-min}})$:

Ratio $R_{\Delta R}$: sum of all neighboring jets / total number of inclusive jets

- For $\Delta R < \pi$ only contributions from ≥ 3 -jet events
- RGE dependence and systematic uncertainties mostly cancel out in the ratio (PDF uncert. $< 3\%$)



used to extract α_s

Strong Coupling Constant



New variable introduced in D0 that characterizes angular correlations of jets and gives an average number of neighboring jets around a reference jet, measured triple differentially: $R_{\Delta R}(p_T, \Delta R, p_{T\text{-nbr-min}})$:


Ratio $R_{\Delta R}$: sum of all neighboring jets / total number of inclusive jets

- For $\Delta R < \pi$ only contributions from ≥ 3 -jet events
- RGE dependence and systematic uncertainties mostly cancel out in the ratio (PDF uncert. $< 3\%$)

$$\alpha_s(M_z) = 0.1191^{+0.0048}_{-0.0071}$$


theor. scale uncertainty dominates

- $\alpha_s(p_T)$ results up to 400 GeV
- $\alpha_s(p_T)$ decreases with p_T as predicted by the RGE
- In agreement with ALEPH, JADE, ZEUS, H1 and world average $\alpha_s(M_z) = 0.1184 \pm 0.0007$




D0 data: 54.5 GeV to 395 GeV coverage


Compilation of α_s results



Results since 2009




V + jet Results




Fixed-order: NLO
LO + Parton Shower
Backgrounds to New Physics

V+jet production


WH production



Single top production



H \rightarrow ZZ \rightarrow 2l2q



Background to top-quark, Higgs, SUSY and other NP productions

- Provide detailed measurements of p_T , mass and angular distributions of vector boson and jets


→ test of fixed order perturbative QCD (MCFM, Blackhat, Rocket, HEJ,...), LO ME+PS predictions in MC event generators (AlpGen, Sherpa, Madgraph,...)

→ testing and tuning of phenomenological models

→ All experiments are heavily involved in such tests

W+jets



- Dominant background to $t\bar{t}$ production, Higgs boson, many non-SM processes
=> extensively studied in all Tevatron and LHC experiments:
jet p_T , H_T , #jets, jet angular, masses, 3rd jet emission prob, etc.
- Good agreement with NLO (Blackhat+Sherpa, HEJ) for most of phase space
(Blackhat: some tension for W+2jet in M_{jj} and high H_T)




M_{jj}

3rd jet emission

H_T




Phys. Rev. D 88, 092001 (2013)





Z+b

- Z+b-jets test of pQCD and b-quark fragmentation, PDF
- Z+b important background for single top, ZH, new phenomena
 - measure ratio with respect to inclusive Z and Z+jet
- Good agreement with NLO predictions (20-25% uncert.) in all experiments


D0	0.0196 ± 0.0012 (stat.) ± 0.0013
(syst.)	
CDF	0.0208 ± 0.0018 (stat) ± 0.0027 (syst.)
MCFM [MSTW2008]	$0.0206^{+0.0022}_{-0.0013}$
ALPGEN+PYTHIA	0.017
SHERPA	0.018




PRD 87, 092010 (2013)



$\sigma(Z+c)/\sigma(Z+jet)$







$\sigma(Z+c)/\sigma(Z+b)$




- D0 differential cross-sections measurements $\sigma_{Z+c-jet}/\sigma_{Z+jet}$ (left) and $\sigma_{Z+c-jet}/\sigma_{Z+b-jet}$ (right) as a function of $p_T(\text{jet})$ ($p_T(\text{jet}) > 20 \text{ GeV}$, $|\eta_{\text{jet}}| < 2.5$).
- Significantly higher than NLO (MCFM) prediction.
- Best agreement is with PYTHIA with $1.7 \times$ enhanced $g \rightarrow cc$ rate.

Photon Production



Direct photons emerge unaltered from the hard subprocess
 → direct colorless probe of the hard scattering dynamics
 → observable: **isolated** photons (typically in $R=0.4$)
 → potential sensitivity to PDFs (gluon!)



+also fragmentation contributions (suppressed by isolation criterion)



Isolated inclusive γ : process composition

Tevatron

LHC





Triple photon+jet differential

Phys. Rev. D 88, 072008 (2013)





- Measurements of diff. cross section in photon p_T in 8 regions:
 $|y^\gamma|$: {0-1, 1.5-2.5}; $|y^{\text{jet}}$: {0-0.8, 0.8-1.6, 1.6-2.4, 2.4-3.2}
- ... with same sign and opposite sign photon and jet rapidities
- Sensitive to parton x from 0.007 to 0.4





Triple photon+jet differential

SS

OS



Central photons
Central jets






Forward photons
Forward jets

Disagreement to NLO at $p_T < 40$ GeV, and $p_T > 70$ GeV with very forward jets.

Photon Pair Production

- Almost irreducible background to $H \rightarrow \gamma\gamma$, other new phenomena => should be understood
- Isolation: $ET_{\text{sum}}[R=0.4] < 2-2.5 \text{ GeV}$ (CDF,D0), $< 4-5 \text{ GeV}$ (Atlas, CMS)
- Min photon p_T varies as 16-20 GeV
- Data are compared with predictions: PYTHIA, SHERPA, DiPhoX, ResBos, NNLO
- 1D cross sections in diphoton Mass, $p_T^{\gamma\gamma}$, $\Delta\phi$, $\cos\theta^*$ and 2D ones ($p_T^{\gamma\gamma}$, $\Delta\phi$, $\cos\theta^*$ in Mass bins)

Diphoton Mass




Small masses: Significant contrib. from $gg \rightarrow \gamma\gamma$ (30-40%), fragmentation

Good agreement with most theories at Mass > 50 GeV, but data are higher than theory up to a factor 1.5-2 at smaller masses.



Photon Pair Production

- Almost irreducible background to $H \rightarrow \gamma\gamma$, other new phenomena, => should be understood
- Isolation: $ET_{\text{sum}}[R=0.4] < 2-2.5 \text{ GeV}$ (CDF,D0), $< 4-5 \text{ GeV}$ (Atlas, CMS)
- Min photon p_T varies as 16-20 GeV
- Data are compared with predictions: PYTHIA, SHERPA, DiPhoX, ResBos, NNLO
- 1D cross sections in diphoton Mass, $p_T^{\gamma\gamma}$, $\Delta\phi$, $\cos\theta^*$ and 2D ones ($p_T^{\gamma\gamma}$, $\Delta\phi$, $\cos\theta^*$ in Mass bins)

$p_T^{\gamma\gamma}$



$\Delta\phi$



D0, PLB 690, 108 (2010)
 D0, PLB 725, 6 (2013)
 CDF, PRL 110, 101801(2013)
 CDF, PRD 84, 052006 (2011)





Significant contrib. from fragmentation photons

NNLO is doing good job, big NNLO/NLO ratio

- None of theories describe the whole phase space: small masses, small $\Delta\phi$ and $\Delta\phi \approx \pi$, moderate $p_T^{\gamma\gamma}$ are most problematic for theories.
- NNLO: good description $p_T^{\gamma\gamma}$, $\Delta\phi$ at CMS, CDF; still should be added the "gg box" HO corrections, resummation (fragm. functions?)

Photon+b

- Disagreement with NLO 5FNS predictions at $p_T > 70$ GeV
- D0 and CDF agree at $p_T > 70$ GeV
- NLO 4FNS describes data within uncertainties.





Tevatron: $q\bar{q} \rightarrow \gamma g (g \rightarrow b\bar{b})$ dominates at $p_T > 80$ GeV
 LHC: $bg \rightarrow b\gamma$ dominates at most p_T

Photon+bb

Submitted to PLB, arXiv:1405.3964




- Measurement of photon+2 b-jet differential cross section vs photon p_T with b-jet $p_T > 20$ GeV, $|y| < 1.5$
- Measurement of ratio $\sigma(\gamma bb)/\sigma(\gamma b)$



- Good agreement with NLO (4FNS) and k_T factorization
- Sherpa underestimates the $\sigma(\gamma bb)/\sigma(\gamma b)$ ratio at low p_T ; Pythia is significantly lower.




Photon+c


PLB 719, 354 (2013)




- Intrinsic charm models predict higher cross sections. BHPS model favored with rise in photon p_T .
- Pythia describes photon+c/photon+b ratio with increased $g \rightarrow cc$ rate (annihilation process)

PLB 719, 354 (2013)







Soft physics



**Phenomenological
nucleon models**

Elastic cross-section

- Measurement of $d\sigma/d|t|$ for $0.25 < |t| < 1.2$ GeV: information on nucleon structure and non-perturbative effects, tests of many phenomenological models.
- Previous measurements: UA4 (546), E710 (1800), CDF (1800)
- Uses a special run with one bunch of p+pbar.



$$d\sigma/dt = A \exp(-b|t|)$$

$$b = 16.86 \pm 0.10 (stat) \pm 0.20 (syst)$$

- Measured fundamental parameter b (tightly related with effective nucleon radius)
- The position of the dip is identified, $|t|=0.6$ GeV²; TOTEM result: dip at $|t|=0.5$ GeV²: diffractive minimum keeps moving to lower $|t|$ values (UA4-->Tevatron-->LHC)




Double parton scattering: Photon+HF+dijet



- Photon: $p_T^\gamma > 26$ GeV, $|\eta| < 1.0$ or $1.5 < |\eta| < 2.5$
- At least 3 jets with $p_T^{jet} > 15$ GeV and $|\eta| < 2.5$
 $15 < p_T^{jet2} < 35$ GeV
- Topology: $\Delta R(\gamma, jet) > 0.7$, $\Delta R(jet, jet) > 1.0$

Case 1: No leading jet flavor requirement
(Inclusive sample)

Case 2: Leading jet Heavy flavor requirement
(HF sample)



Check dependence on initial parton flavor!

$$\sigma_{DP} = \sigma_A \sigma_B / \sigma_{eff}$$

σ_{eff} is directly related with parton spatial density.


Data	1Vtx	2Vtx
Sample		
inclusive	218686	269445
HF	5004	5811

Fractions of Double Parton events




DP event fraction is found by maximum likelihood fitting Single Parton event model (Sherpa) and Double Parton signal event model (MixDP) to data.

Inclusive sample



Heavy Flavor sample




DP fractions

γ +HF+dijet	γ +3 jet
0.171 ± 0.020	0.202 ± 0.007

Effective cross section

Phys.Rev.D89, 072006 (2014), arXiv:1402.1550

- Having measured number of DP events and corresponding acceptances and efficiencies one can calculate σ_{eff} for both final states.
- Measured σ_{eff} is in agreement with all Tevatron and LHC measurements, but the new values is more precise.
- No dependence of σ_{eff} on initial quark flavor has been observed.





Final state	γ +HF+dijet	γ +3 jet
σ_{eff} (mb)	14.6 ± 3.26	12.7 ± 1.32

Double J/psi production



- Dominant production channel: $gg \rightarrow J/\psi J/\psi$
- Signal: prompt direct J/ψ (S-wave) and (P-waves) $^m_{1c}$ and $^m_{2c}, ^m_{1(2)c} \rightarrow J/\psi + \gamma$
- Background: non-prompt B-hadron decays, non-resonant DY , π/K decays.
- Single and Double parton scatterings may contribute
 \Rightarrow Test of σ_{eff} energy dependence: from high energies to 4-5 GeV,
 with gg initial state only

Prediction for the Tevatron at $p_T(J/\psi) > 4$ GeV, $|\eta| < 0.6$:
 expected DP fraction is $\sim 15\%$



Single Parton and Double Parton contributions

- We measure the Double J/ψ production cross section for Double Parton and Single Parton scatterings separately. To discriminate between the two mechanisms, we use $\Delta\eta(J/\psi, J/\psi)$ difference.

- Contributions from double non-prompt, prompt+non-prompt and accidental backgrounds are subtracted from data
 => data should contain just prompt SP and DP events.


SP template: DJ events simulated with Herwig++ /DJPsiFDC
 DP template: Pythia-8 or data-like DP model.

Systematics: fit and variation between the 2+2 models; prompt+non-prompt origin (either 100% SP- or DP-like). DP double non-prompt is highly suppressed to 0.7-2 fb.

$$\sigma_{eff} = \frac{\sigma(J/\psi)^2}{\sigma(J/\psi J/\psi)}$$

$$\sigma_{eff} = 5.0 \pm 0.5 (stat) \pm 2.7 (syst) mb$$

DP dominates at $|\Delta\eta(J/\psi J/\psi)| > 2$



$$f_{SP} = 0.70 \pm 0.11, f_{DP} = 0.30 \pm 0.10$$




Summary

- Good consistency and complementarity for most of experimental data
- Current level of understanding jet ID, systematics and jet energy scale leads in many cases to **experimental uncertainties similar or lower than theoretical uncertainties.**
- **Jet results: Precision measurement of fundamental observables.**
=> sensitivity to PDF sets, strongest constraint on gluon PDF, extraction of α_s and test of its running up to 400 GeV, detailed studies of the effect of different jet algorithms, study of jet substructure, limits on many NP models.
- **Z/W results:** extensive tests of pQCD and MC models; in most cases, a triumph of NLO and ME-PS MC predictions.
- **Photon results:** test of fixed order NLO, resummation, fragmentation. Theory should be better understood. First NNLO results look very promising.
- **UE/DP events:** improving phenomenological models, good knowledge is required in multijet studies/searches.

BACK-UP SLIDES

Pion cloud model

- For details please see e.g. PRD80:114029,2009, PRD83:054012,2
 - In this model, there can be interactions of gluons and quarks in the proton "core" with soft pions in the "cloud".
- The "bare" parton can make transition to a virtual state containing a pion, $p \rightarrow n\pi^+$ (more likely), $p \rightarrow \Delta^{++}\pi^-$ (excess of π^+ vs π^- , $\Rightarrow |\bar{d}-\bar{u}| > 0$ for the "sea" quarks). The pion is a "slow parton", with a momentum $y = M_\pi / M_p$




Due to these interactions, u and d quarks size grows more rapidly than gluonic radius

$$\langle b^2 \rangle_f = \frac{\int d^2b b^2 [f(x,b)_{\text{core}} + \Theta(b > b_0) f(x,b)_{\text{chiral}}]}{f(x)}$$

$$\equiv \langle b^2 \rangle_{f,\text{core}} + \langle b^2 \rangle_{f,\text{chiral}} \quad (56)$$

In numbers, for $x \sim 0.01$, we get about 30% larger $\langle b^2 \rangle_{q+qbar}$ than $\langle b^2 \rangle_g$

Assumption made: transverse sizes of quarks and gluons in the core are the same (is it true?).




Pion cloud model: GPD and experiment

In GPD formalism, the transverse quark/gluon sizes are related to the corresp. GPD:

$$\langle b^2 \rangle_f(x) \equiv \frac{\int d^2b b^2 f(x,b)}{f(x)} \quad \leftarrow \text{GPD} \quad (f = q, \bar{q}, g)$$

$$\langle b^2 \rangle_g = 4 \frac{\partial}{\partial t} \left[\frac{d\sigma/dt(t)}{d\sigma/dt(0)} \right]_{t=0}^{1/2} \quad (d\sigma/dt)^{\gamma N \rightarrow J/\psi + N} \propto \exp(B_{J/\psi} t)$$

$$\langle b^2 \rangle_g = 2B_{J/\psi}$$



t-dependence of J/ψ photoproduction cross section is sensitive to the transverse gluon size. H1, ZEUS data:

$$B_{J/\psi} \approx 4.1 - 4.6 \text{ GeV}^{-2} \text{ or } \langle b^2 \rangle_g = 0.32 - 0.35 \text{ fm}^2$$

t-dependence of deeply-virtual Compton scattering cross section is sensitive to the transverse quark size. H1 data:

$$B_\gamma \approx 5.2 - 5.8 \text{ GeV}^{-2} \text{ or } \langle b^2 \rangle_{q+qbar} = 0.42 - 0.46 \text{ fm}^2$$

See PRD80:114029,2009, C.Weiss, DIS2011.


Dijet mass: searches for new physics

PRD 79, 112002 (2009)

Dijet mass tests pQCD but also sensitive to presence of new physics, resonances decaying to two jets

=> Use uncorrected jet data to maximize sensitivity to resonances

No significant evidence for resonant structure has been observed, so set limits





↓

Observed mass exclusion range	Model description
260-870 GeV/c^2	Excited quark $\rightarrow qg$ ($f=f'=f_s=1$)
260-1100 GeV/c^2	ρ_{T8} techni-rho
260-1250 GeV/c^2	Axigluon/coloron
290-630 GeV/c^2	E_6 diquark
280-840 GeV/c^2	W' (SM couplings)
320-740 GeV/c^2	Z' (SM couplings)

D0 dijet χ : limits on q-compositeness, Extra Dim.: PRL 103, 191803 (2009)

“Tevatron” to the LHC




- Shows the “transAVE” **charged particle density** as defined by the leading charged particle, PTmax, as a function of PTmax at sqrt(s)=300 GeV, 900 GeV, 1.96 TeV, and 7 TeV compared with the CMS PYTHIA 8 tune CUETP8S1-CTEQ6L.

- Shows the “transAVE” **charged PTsum density** as defined by the leading charged particle, PTmax, as a function of PTmax at sqrt(s)=300 GeV, 900 GeV, 1.96 TeV, and 7 TeV compared with the CMS PYTHIA 8 tune CUETP8S1-CTEQ6L.

What we are learning should allow for a deeper understanding of MPI which will result in more precise predictions at the future LHC energies of 13 & 14 TeV

“Tevatron” to the LHC



Having determined the parameters of an MPI model, one can make an unambiguous prediction of σ_{eff} . In PYTHIA 8 σ_{eff} depends primarily on the matter overlap function, which for bProfile = 3 is determined by the exponential shape parameter, expPow, and the MPI cross section determined by p_{TO} and the PDF.


- Shows the “transAVE” as defined by the leading charged particle, PTmax, as a function of PTmax at 300 GeV, 900 GeV, 1.96 TeV, and 7 TeV compared with the CMS PYTHIA 8 tune CUETP8S1-CTEQ6L.
- Shows the “transAVE” as defined by the leading charged particle, PTmax, as a function of PTmax at 300 GeV, 900 GeV, 1.96 TeV, and 7 TeV compared with the CMS PYTHIA 8 tune CUETP8S1-CTEQ6L.

What we are learning should allow for a deeper understanding of MPI which will result in more precise predictions at the future LHC energies of 13 & 14 TeV


Sigma-Effective

Experiment, Final state (Year)

σ_{eff} measurements




New D0 values



- Shows the σ_{eff} values calculated from the PYTHIA 8 Monash and CMS tune CUETP8S1-CTEQ6L.

The σ_{eff} predicted from the PYTHIA 8 UE tunes is slightly larger than the direct measurements!

Sigma-Effective




M. H. Seymour^a A. Siódmok^a

^a Consortium for Fundamental Physics, School of Physics and Astronomy, The University of Manchester, Manchester, M13 9PL, U.K.

E-mail: michael.seymour@manchester.ac.uk,
andrzej.siodmok@manchester.ac.uk

ABSTRACT: We review the modelling of multiple interactions in the event generator HERWIG++ and study implications of recent tuning efforts to Tevatron and LHC data.

It is often said that measurements of the effective cross section for double-parton scattering, σ_{eff} , are in contradiction with models of the final state of multi-parton interactions, but we show that the HERWIG++ model is consistent with both and gives stable predictions for underlying event observables at 14 TeV.



Parton spatial density and σ_{eff}

Double parton cross section

$$\sigma_{\text{dp}} = \sum_{q/g} \int \frac{\sigma_{12}\sigma_{34}}{2\sigma_{\text{eff}}} D_p(x_1, x_3) D_{\bar{p}}(x_2, x_4) dx_1 dx_2 dx_3 dx_4$$


Effective cross section σ_{eff} is directly related with parton spatial density:

$$\sigma_{\text{eff}} = \left[\int d^2\beta [F(\beta)]^2 \right]^{-1}$$

β is impact parameter


$$F(\beta) = \int f(b) f(b-\beta) d^2b$$

where $f(b)$ is the density of partons in transverse space.
 => Having σ_{eff} measured we can estimate $f(b)$



Double parton scattering

Inclusive Jets: Tevatron vs. LHC



PDF sensitivity:

→ compare jet cross section at fixed
 $x_T = 2 p_T / \sqrt{s}$

Tevatron (ppbar)

>100x higher cross section @ all x_T
>200x higher cross section @ $x_T > 0.5$

LHC (pp)

- need more than 2400 fb^{-1} luminosity to improve Tevatron@ 12 fb^{-1}
- more high-x gluon contributions
- but more steeply falling cross sect. at highest p_T (=larger uncertainties)


W+c

PRL 110, 071801 (2013)




Sensitive to s-quark PDF: 90% s, 10% d

signal: OS \gg SS
 backgrounds: OS \sim SS
 (subtracted in the diff. 'OS-SS')



Lepton p_T in jet



- Jet $p_T > 20$ GeV, $|\eta| < 1.5$
- 5.7σ CL for W+single c-jet

• $\sigma(W+c) \cdot Br(W \rightarrow l\nu)$:

CDF Data: $13.6^{+3.4}_{-3.1}$ pb



QCD NLO: 11.4 ± 1.3 pb

- Also measured $|V_{cs}| = 1.08 \pm 0.16$

Good agreement
 data/theory

Measurements of $\sigma(W+D^*)/\sigma(W)$



Preliminary



- CDF data for the differential rates of cross-section ratio $\sigma(W + D^*)/\sigma(W)$ as a function of $p_T(D^*)$, as measured by in the $W \rightarrow e\nu$ (left) and $W \rightarrow \mu\nu$ (right) decay channels. D^* is fully reconstructed at the track level [$D^*(2010) \rightarrow D0(\rightarrow K\pi)\pi_s$]

Measurements of $\sigma(V+D^*)/\sigma(V)$

Preliminary



Production process ($p_T(D^*) > 3 \text{ GeV}/c$)	CDF Run II Preliminary $\int \mathcal{L} dt = 9.7 \text{ fb}^{-1}$ $\sigma(V + D^*)/\sigma(V)$ (%) $\pm(\text{stat}) \pm(\text{syst})$	Pythia 6.2.16 (CTEQ5L) $\sigma(V + D^*)/\sigma(V)$ (%) $\pm(\text{pdf unc})$
$W(\rightarrow e\nu) + D^*$	$1.74 \pm 0.21 \pm 0.17$	1.77 ± 0.07
$W(\rightarrow \mu\nu) + D^*$	$1.75 \pm 0.17 \pm 0.05$	1.77 ± 0.07
Combined results: $W(\rightarrow e\nu/\mu\nu) + D^*$	$1.75 \pm 0.13 \pm 0.09$	1.77 ± 0.07
$Z(\rightarrow ee) + D^*$	$1.0 \pm 0.6 \pm 0.2$	1.36 ± 0.05
$Z(\rightarrow \mu\mu) + D^*$	$1.8 \pm 0.5 \pm 0.2$	1.36 ± 0.05
Combined results: $Z(\rightarrow ee/\mu\mu) + D^*$	$1.5 \pm 0.4 \pm 0.2$	1.36 ± 0.05

- CDF data for the differential rates of cross-section ratio $\sigma(W + D^*)/\sigma(W)$ as a function of $p_T(D^*)$, as measured by in the $W \rightarrow e\nu$ (left) and $W \rightarrow \mu\nu$ (right) decay channels.
 D^* is fully reconstructed at the track level [$D^*(2010) \rightarrow D0(\rightarrow K\pi)\pi_s$]
- The measurements show good agreement with PYTHIA 6.2 Tune A with in all bins.


W/Z+Upsilon Search

Preliminary

- CDF search for the production of the Upsilon (1S) meson in association with a vector boson.
- 9.7 fb⁻¹ data set



Observe one Upsilon + W candidate over an expected background of 1.2 ± 0.5 events, and one Upsilon + Z candidate over an expected background of 0.1 ± 0.1 events.



	$\Upsilon + W \rightarrow e\nu$	$\Upsilon + W \rightarrow \mu\nu$	$\Upsilon + W \rightarrow l\nu$	$\Upsilon + Z \rightarrow ee$	$\Upsilon + Z \rightarrow \mu\mu$	$\Upsilon + Z \rightarrow ll$
N_{sig}	0.019±0.004	0.014±0.003	0.034±0.006	0.0048±0.0009	0.0037±0.0007	0.0084±0.0016
N_{bg} (fake Υ)	0.7±0.4	0.4±0.3	1.1±0.5	0.07±0.07	0.04±0.04	0.1±0.1
N_{bg} (fake W/Z)	0.06±0.04	negl.	0.06±0.04	negl.	negl.	negl.
N_{bg} ($\Upsilon + Z$)	0.0006±0.0001	0.0033±0.0006	0.0039±0.0007			
N_{bg} (total)	0.8±0.4	0.4±0.3	1.2±0.5	0.07±0.07	0.04±0.04	0.1±0.1
N_{obs}	0	1	1	0	1	1



95% C.L. Cross Section Limits

	$\Upsilon + W$	$\Upsilon + Z$
expected limit (pb)	5.5	13
observed limit (pb)	5.5	20

Strange particle production



- Strangeness production is used for refining phenomenological models and parameters of the Monte Carlo models.
- Enhanced production of the strange particle has been frequently suggested as a manifestation of the formation of quark-gluon plasma.
- Measured production cross section of $\Lambda(uds)$, $\Xi^\pm(.ss)$ and $\Omega^\pm(sss)$



Cross sections depend on the number of strange quarks, however very similar p_T slopes indicate an universality in particle production.

Exclusive di-photons



Search for exclusive $\gamma\gamma$ production via $p\bar{p} \rightarrow p + \gamma\gamma + \bar{p}$ and compare to theory

Motivation: intrinsically interesting QCD process;

tightly related with excl. Higgs boson production $p\bar{p} \rightarrow p + H + \bar{p}$




Features:

- a) proton and antiproton emerge intact with no hadrons produced
- b) .. have $p_T < 1$ GeV, having emitted a pair of gluons (in CS mode)

Event selection:

- two central photons with $p_T > 2.5$ GeV
- No other activity in the detector

Background: irreducible ($q\bar{q} \rightarrow \gamma\gamma$) $< 5\%$
 reducible ($\pi^0\pi^0, \eta\eta$) are $< 16\%$




PRL 108, 081801 (2012)

Regge theory: diffractive scattering via pomeron exchange

Data: $2.48^{+0.40}_{-0.35} (stat)^{+0.40}_{-0.51} (syst)$

Good agreement with theory



Exclusive di-photons



PRL 108, 081801 (2012)

Search for exclusive $\gamma\gamma$ production via $p\bar{p} \rightarrow p + \gamma\gamma + \bar{p}$ and compare to theory

Motivation: intrinsically interesting QCD process;

tightly related with excl. Higgs boson production $p\bar{p} \rightarrow p + H + \bar{p}$





Features:

- a) proton and antiproton emerge intact with no hadrons produced
- b) .. have $p_T < 1$ GeV, having emitted a pair of gluons (in CS mode)

Event selection:

- two central photons with $p_T > 2.5$ GeV
- No other activity in the detector

Background: irreducible ($q\bar{q} \rightarrow \gamma\gamma$) $< 5\%$
 reducible ($\pi^0\pi^0, \eta\eta$) are $< x\%$




Other exclusive diffractive productions:

- dijets, D0: PLB 705, 193 (2011)
- dijets, CDF: PRD 86, 032009 (2012)
- W/Z, CDF: PRD82, 112004 (2010)
- Charmonium, CDF: PRL, 102, 242001 (2009)
- e+e-, CDF: PRL 98, 112001 (2007)

W+b


- Tests of pQCD
- Very important background to SUSY and Higgs boson searches
- Measured total *exclusive* W+b cross sections, up to two jets in the final state with jet $p_T > 20$ (25) GeV and $|y| < 2.0$ (2.1) at CDF (Atlas)



~90% (50%)@ TeV(LHC)

CDF

PRL 104, 131801(2010)



$\sigma(W+b\text{-jets}) \cdot \text{BR}(W \rightarrow lv)$ [pb]:

Data: $2.74 \pm 0.27(\text{stat}) \pm 0.42(\text{syst})$

NLO: 1.20 ± 0.14


Data/NLO

Difference: $\sim 3\sigma$

about factor 2

ATLAS

PLB 707, 418(2012)



W+1b:
agreement

$10.2 \pm 1.9(\text{stat}) \pm 1.6(\text{syst})$
 $4.8_{-0.7}^{+1.2}(\text{scale})_{-0.0}^{+0.3}(\text{PDF})_{-0.2}^{+0.3}(m_b) \pm 0.3(\text{np.corr})$



$\sim 1.5\sigma$

=> Should be cross-checked in at least one more measurement, preferably *inclusive* and *differential*

Heavy Flavor Jet Identification

➔ B and C hadrons have a relatively long lifetime (~ 1 ps) and travel ~ 100 - $500\mu\text{m}$ before decay with large masses 2-5 GeV

- Tracks displaced from primary vertex with large impact parameters




➔ Heavy flavor tagging exploits these characteristics of the tracks to create a discriminant used to enrich sample with HF-jets


Heavy Flavor Jet Identification

➔ B and C hadrons have a relatively long lifetime (~ 1 ps) and travel ~ 100 - $500\mu\text{m}$ before decay with large masses 2-5 GeV

- Tracks displaced from primary vertex with large impact parameters



Inv. Mass of Secondary Vertex



➔ Heavy flavor tagging exploits these characteristics of the tracks to create a discriminant used to enrich sample with HF-jets

Z/W+jets production

Use leptonic Z/W decays as most precise probe of QCD

- high Q^2 ($\sim M_Z$ or M_W)
- very small backgrounds, right down to very small p_T !

Concentrate on high p_T final states

- regime of perturbative QCD (and NP)

Theory predictions:

pQCD (+ corrections for underlying event & hadronization):

- LO Z(W) + 1 - 6 partons
- NLO Z(W) + 1, 2 (MCFM)
- NLO W+3[4] (Rocket+MCFM, Blackhat+SHERPA)

Event generators:

- LO $2 \rightarrow 1, 2$ + parton shower
 - PYTHIA, HERWIG
- LO $2 \rightarrow 1-6$ + (vetoed) parton shower
 - ALPGEN (MLM ME-PS matching),
 - SHERPA (CKKW ME-PS matching)

These generators are the main Tevatron and LHC tools

- but, leading order \rightarrow large uncertainties
- must be tuned to data!

