

# Measurement of cross-sections of yttrium ( $n, xn$ ) threshold reactions by means of gamma spectroscopy

XXII ISHEPP

P. Chudoba <sup>1,2</sup>, S. Kilim <sup>3</sup>, Polish <sup>3</sup> and Czech <sup>1</sup> group

<sup>1</sup>Nuclear Physics Institute of the ASCR, v. v. i., Rez 130, 250 68 Rez, Czech Republic

<sup>2</sup>Charles University in Prague, Faculty of Mathematics and Physics, Ke Karlovu 3, 121 16 Praha 2, Czech Republic

<sup>3</sup>National Centre for Nuclear Research, Otwock-Świerk 05-400, Poland

16.9.2014

# Outline

- 1 Motivation of measurement
- 2 Cross-sections measurement
- 3 Experimental arrangement
- 4 Cross-section results

# Motivation of measurement

Necessity of fast neutron field monitoring in facilities like

- Accelerator driven systems (ADS)
- Neutron spallation sources
- Future fusion reactors and fast reactors

# Why yttrium?

- Products of (n,xn) reactions on yttrium are easily identifiable
- Half-lives of the products have good length for  $\gamma$ -spectrometry
- $\gamma$  transitions are intensive enough for detection and are well separated from each other

Reaction	$E_{thr}$ [MeV]	$T_{\frac{1}{2}}$	$E_{\gamma}$ [keV]	$I_{\gamma}$ [%]
$^{89}\text{Y}(n,2n)^{88}\text{Y}$	11.6	106.626 d	898.042	93.683
			1836.063	99.24
$^{89}\text{Y}(n,3n)^{87}\text{Y}$	21.1	79.8 h	388.531	82.2
			484.805	89.845
$^{89}\text{Y}(n,3n)^{87m}\text{Y}$	21.6	13.37 h	380.79	78.055

# $\gamma$ -spectroscopic methods


- With accurate knowledge of isotopic composition, it is possible to measure cross-section
- With known cross-section the integral of the beam is possible to determine
- Investigated reactions:  $^{89}\text{Y}(n,2n)^{88}\text{Y}$ ,  $^{89}\text{Y}(n,3n)^{87}\text{Y}$
- Equations for determining of cross-section in case of simple decay

$$N_{\text{yield}} = \frac{S_{\text{peak}} \cdot C_{\text{abs}}(E)}{I_{\gamma} \cdot \varepsilon_p(E) \cdot \text{COI}(E) \cdot C_{\text{area}}} \frac{t_{\text{real}}}{t_{\text{live}}} \frac{e^{\lambda \cdot t_0}}{1 - e^{-\lambda \cdot t_{\text{real}}}} \frac{\lambda \cdot t_{\text{irr}}}{1 - e^{-\lambda \cdot t_{\text{irr}}}},$$

$$\sigma = \frac{N_{\text{yield}} \cdot S \cdot A \cdot B_a}{N_n \cdot N_A \cdot m}.$$


# Self-absorption correction - $C_{abs}(E)$

$$C_{abs} = \frac{\int_0^D \frac{I_0}{D} dx}{\int_0^D \frac{I_0}{D} e^{-\mu \cdot x} dx} = \frac{\mu \cdot D}{1 - e^{-\mu \cdot D}}$$





# Square emitter correction - $C_{area}$

$$C_{area} = \frac{\epsilon_{foil}}{\epsilon_{point}}$$




# Detector efficiency simulations






# Comparison between experiment and simulation




# Comparison between experiment and simulation




# Decay of $^{87}\text{Y}$ and $^{87m}\text{Y}$

In case of decay of  $^{87}\text{Y}$  and  $^{87m}\text{Y}$  the shape is a little more complicated.



# Correction for decay during cooling and irradiation

- In case of  $^{89}\text{Y}(n,3n)^{87}\text{Y}$  reaction the ground and isomeric states are produced simultaneously
- For determination of  $^{87}\text{Y}$  ground state production it is necessary to involve  $^{87m}\text{Y}$  decay to the ground state.





$$N_{g0} = \frac{S_{peak} \cdot C_{abs}(E) \cdot B_a \cdot e^{\lambda_g \cdot t_0} \cdot t_{real}}{I_\gamma \cdot \epsilon_p(E) \cdot COI(E) \cdot C_{area} \cdot t_{live}} \cdot \frac{1}{(1 - e^{-\lambda \cdot t_{real}})} + \frac{\lambda_m \cdot N_{m0}}{\lambda_g - \lambda_m} \left(1 - e^{(\lambda_g - \lambda_m) \cdot t_0}\right)$$

$$C_{irr} = \frac{P_g \cdot t_{irr}}{N_{g0}} = \frac{\lambda_g \cdot t_{irr}}{1 - e^{-\lambda_g \cdot t_{irr}}} - \frac{N_{yield,m}}{N_{g0} \cdot (1 - e^{-\lambda_g \cdot t_{irr}})} \left(1 - \frac{\lambda_m \cdot e^{-\lambda_g \cdot t_{irr}} - \lambda_g \cdot e^{-\lambda_m \cdot t_{irr}}}{\lambda_m - \lambda_g}\right)$$



$$N_{yield,g} = N_{g0} \cdot C_{irr}$$

# Fast neutron source

- Source based on reaction  ${}^7\text{Li}(p,n){}^7\text{Be}$
- Neutron energy range 10-37 MeV
- Source intensity  $\sim 10^8 \text{ cm}^{-2}\cdot\text{s}^{-1}$
- Neutron spectrum is obtained by an MCNPX simulation



# Model of the neutron source with samples





Two yttrium samples for each irradiation, each irradiated with gold foil:

- YN -  $25 \times 25 \times 0.64$  mm - solid foil, distance 123 mm
- YO -  $\varnothing 9 \times 1.5$  mm - pill, distance 103 mm

# Background subtraction

- Neutron source is not monoenergetic
- Neutron background subtraction based on folding of neutron spectrum and cross-section


$$C_{bgr} = \frac{\int_{Peak} \sigma(E) \cdot N(E) dE}{\int_{Spectrum} \sigma(E) \cdot N(E) dE} \longrightarrow C_{bgr} = \frac{\sum_{i \in Peak} \sigma_i \cdot N_i}{\sum_i \sigma_i \cdot N_i}$$



# Measurement equipment


## HPGe spectrometer CAN35

- Relative efficiency 35%
- Calibration points from 53 to 1836 keV
- Calibrated positions 3, 4, 5, 6, 7, 10, 12, 15, 17 cm
- Complete shielding









# Cross-section of $^{89}\text{Y}(n,x\text{n})$ reaction



# Cross-section of $^{197}\text{Au}(n,xn)$ reaction



!Preliminary!

# Thank you for your attention

The experiment was supported by ERINDA<sup>1</sup> and CANAM<sup>2</sup>.

---

<sup>1</sup><http://www.erinda.org>

<sup>2</sup><http://canam.ujf.cas.cz>