RENORMDYNAMICS AND PRODUCTION OF FRACTALS, FLUCTONS, SOLITONS, STRINGS, UNPARTICLES ...

Nugzar Makhaldiani

JINR Dubna, mnv@jinr.ru

Talk at the XXIII International Baldin Seminar on High Energy Physics Problems
"Relativistic Nuclear Physics and Quantum Chromodynamics"
September 19 - 24, 2016 JINR Dubna, Russia

"I do not know what I may appear to the world;
but to myself I seem to have been only like a boy
playing on the sea-shore, and
diverting myself in now and then
finding a smoother pebble or
a prettier shell than ordinary,
whilst the great ocean of truth
lay all undiscovered before me."
(Isaac Newton, In Brewster,
Memoirs of Newton (1855), vol II, Ch. 27)

"... all things physical are information-
theoretic in origin ..."
John A. Wheeler

Why supersymmetry is
So universal?

New Physics

We say that we find **New Physics** (NP) when either we find a phenomenon which is forbidden by SM in principal - this is the qualitative level of NP - or we find a significant deviation between precision calculations in SM of an observable quantity and a corresponding experimental value.
In 1900, the British physicist Lord Kelvin is said to have pronounced: "There is nothing new to be discovered in physics now. All that remains is more and more precise measurement." Within three decades, quantum mechanics and Einstein's theory of relativity had revolutionized the field. Today, no physicist would dare assert that our physical knowledge of the universe is near completion. To the contrary, each new discovery seems to unlock a Pandora's box of even bigger, even deeper physics questions.

In the Universe, matter has manly two geometric structures, homogeneous, [Weinberg, 1972] and hierarchical, [Okun, 1991] .
The homogeneous structures are naturally described by real numbers with an infinite number of digits in the fractional part and usual archimedean metrics. The hierarchical structures are described with p-adic numbers with an infinite number of digits in the integer part and non-archimedean metrics, [Koblitz, 1977].

A discrete, finite, regularized, version of the homogenous structures are homogeneous lattices with constant steps and distance rising as arithmetic progression. The discrete version of the hierarchical structures is hierarchical lattice-tree with scale rising in geometric progression.

There is an opinion that present day theoretical physics needs (almost) all mathematics, and the progress of modern mathematics is stimulated by fundamental problems of theoretical physics.

In QFT existence of a given theory means, that we can control its behavior at some scales (short or large distances) by renormalization theory [Collins, 1984].
If the theory exists, than we want to solve it, which means to determine what happens on other (large or short) scales. This is the problem (and content) of Renormdynamics.
The result of the Renormdynamics, the solution of its discrete or continual motion equations, is the effective QFT on a given scale (different from the initial one).
We can invent scale variable λ and consider QFT on $D+1+1$ dimensional space-time-scale. For the scale variable $\lambda \in (0,1]$ it is natural to consider q-discretization, $0<q<1,\ \lambda_n = q^n,\ n=0,1,2,\ ...$ and p - adic, nonarchimedian metric, with $q^{-1}=p$ - prime integer number.
The field variable $\varphi(x,t,\lambda)$ is complex function of the real, x, t, and p - adic, λ, variables. The solution of the UV renormdynamic problem means, to find evolution from finite to small scales with respect to the scale time $\tau = \ln \lambda/\lambda_0 \in (0,-\infty)$. Solution of the IR renormdynamic problem means to find evolution from finite to the large scales, $\tau = \ln \lambda/\lambda_0 \in (0,\infty)$.

We will call RDF functions $g_n = f_n(t)$ which are solutions of the RD motion equations

$$\dot{g}_n = \beta_n(g), 1 \leq n \leq N. \tag{1}$$

In the simplest case of one coupling constant the function $g = f(t)$ is constant, $g = g_c$ when $\beta(g_c) = 0$, or is invertible (monotone). Indeed,

$$\dot{g} = f'(t) = f'(f^{-1}(g)) = \beta(g). \tag{2}$$

Each monotone interval ends by UV and IR fixed points and describes corresponding phase of the system.
Note that the simplest case of the classical dynamics, the Hamiltonian system with one degree of freedom, is already two dimensional, so we have no analog of one charge renormdynamics.

The regular Hamiltonian systems of the classical mechanics are defined on the even dimensional phase space, so there is no analog of the three dimensional renormdynamics for the coupling constants of the SM. The fixed points of renormdynamics belong to a set of solutions of the polynomial system of equations $\beta_n(g) = 0, 1 \leq n \leq N$, in the perturbative renormdynamics. Describing the solutions is the task of contemporary algebraic and computational geometry.

The quantitative values and qualitative content of the given field theory depend on the scale (parameter, e.g. $\mu-$renormalization point, $g = g(\mu),\ A = A(\mu)$). In QCD e.g. the effective action has the following form:

$$S(\mu) = \frac{1}{g^2(\mu)} \int d^D x \mathcal{L}(A(\mu)), \tag{3}$$

variation with respect to the change of scale gives

$$\delta S = -2\frac{\beta(g)}{g\mu}\delta\mu S + \frac{1}{g^2}\int d^D x \frac{\delta\mathcal{L}}{\delta A}\delta A \tag{4}$$

and the following two statements are equivalent:

$$\delta S = 0,\ \beta(g) = 0 \Leftrightarrow \delta S = 0,\ \frac{\delta\mathcal{L}}{\delta A} = 0. \tag{5}$$

So, from renorminvariance of the effective action follows that at the conformal symmetric points, the motion equations for fields are satisfied. Generalization for the several coupling constants and other models is obvious.

In string theory, the connection between conformal invariance of the effective theory on the parametric world sheet and the motion equations of the fields on the embedding space is well known [Ketov, 2000]. A more recent topic in this direction is AdS/CFT Duality [Maldacena, 1988]. In this approach for QCD coupling constant the following expression was obtained [Brodsky, de Tèramond, Deur, 2010]

$$\alpha_{AdS}(Q^2) = \alpha(0) e^{-Q^2/4k^2}. \tag{6}$$

A corresponding β-function is

$$\beta(\alpha_{AdS}) = \frac{d\alpha_{AdS}}{\ln Q^2} = -\frac{Q^2}{4k^2}\alpha_{AdS}(Q^2) = \alpha_{AdS}(Q^2) \ln \frac{\alpha_{AdS}(Q^2)}{\alpha(0)} \leq 0 \tag{7}$$

So, this renormdynamics of QCD interpolates between the IR fixed point $\alpha(0)$, which we take as $\alpha(0) = 2$, and the UV fixed point $\alpha(\infty) = 0$.

For the QCD running coupling considered in [Diakonov, 2003]

$$\alpha(q^2) = \frac{4\pi}{9\ln(\frac{q^2+m_g^2}{\Lambda^2})}, \tag{8}$$

where $m_g = 0.88GeV,\ \Lambda = 0.28GeV$, the $\beta-$function of renormdynamics is

$$\beta(\alpha) = -\frac{\alpha^2}{k}(1 - c\exp(-\frac{k}{\alpha})),$$
$$k = \frac{4\pi}{9} = 1.40,\ c = \frac{m_g^2}{\Lambda^2} = (3.143)^2 = 9.88, \tag{9}$$

for a nontrivial (IR) fixed point we have

$$\alpha_{IR} = k/\ln c = 0.61 \tag{10}$$

For $\alpha(m) = 2$, at valence quark scale m we predict the gluon (or valence quark) mass as

$$m_g = \Lambda e^{\frac{k}{2\alpha(m)}} = 1.42\Lambda = m_N/3,\ \Lambda = 220MeV. \tag{11}$$

Valence Quark Coupling Constant

It is nice to have a nonperturbative $\beta-$function like (9), but it is more important to see which kind of nonperturbative corrections we need to have a phenomenological coupling constant dynamics.
It was noted [Voloshin, Ter-Martyrosian, 1984] that in valence quark parametrization $\alpha_s(m) = 2$, at a valence quark scale m.

In the 1870's G.J. Stoney [Stoney, 1881], the physicist who coined the term "electron" and measured the value of elementary charge e, introduced as universal units of Nature for L, T, M :

$$l_S = \frac{e}{c^2}\sqrt{G},\ t_S = \frac{e}{c^3}\sqrt{G},\ m_S = \frac{e}{\sqrt{G}} \tag{12}$$

M. Planck introduced [Planck, 1899] as universal units of Nature for L, T, M:

$$m_P = \sqrt{\frac{hc}{G}} = \frac{m_S}{\sqrt{\alpha}},\ l_P = \frac{h}{cm_P} = \frac{l_S}{\sqrt{\alpha}} = 11.7 l_S,\ t_P = \frac{l_P}{c} = \frac{t_S}{\sqrt{\alpha}} \tag{13}$$

Stoney's fundamental constants are more fundamental just because they are less than Planck's constants :) Due to the value of $\alpha^{-1} = 137$, we can consider relativity theory and quantum mechanics as deformations of the classical mechanics when deformation parameter $c = 137$ (in units $e = 1, \hbar = 1$) and $\hbar = 137$ (in units $e = 1, c = 1$), correspondingly. These deformations have an analytic sense of p-adic convergent series. The number 137 has a very interesting geometric sense,

$$137 = 11^2 + 4^2, \tag{14}$$

so, $\sqrt{137}$ is the hypotenuse length of a triangle with other sides of lengths 11 and 4.

Base of the Babylonians Number System

The Babylonians used a base 60 number system which is still used for measuring time - 60 seconds in a minute, 60 minutes in an hour - and for measuring angle - 360 degrees in a full turn. The base 60 number system has its origin in the ration of the Sumerian mina (m) and Akkadian shekel (s), $m/s \simeq 60 = 3 \cdot 4 \cdot 5$.
We also can consider base 137 system for fundamental theories.

Base of the Fundamental Number System

For the nuclear physics strong coupling phenomena description we may take as a base $p = 13$.
For the hadronic physics, valence scale QCD, and graphen strong coupling phenomena description we may take as a base $p = 2$.
For the weak coupling physics SM m_Z scale and MSSM unification scale phenomena description we may take as a base $p = 29$.

Number of the Fundamental Constants

There are different opinions about the number of fundamental constants [Duff, Okun, Veneziano, 2001].
According to Okun, there are three fundamental dimensionful constants in Nature: Planck's constant, $\hbar$; the velocity of light, c; and Newton's constant, G.
According to Veneziano, there are only two: the string length L_s and c.
According to Duff, there are not fundamental constants at all.
Usually $L_s = l_p$, so, the fundamental area is $L_s^2 = 137 l_s^2$.
The value $s_s = l_s^2$– Stoney area, is more like on a fundamental area :)

Scale Dependent Number of Fundamental Constants

In mathematics we have two kind of structures, discrete and continuous one. If a physical quantity has discrete values, it might have no dimension. If the values are continuous - the quantity might have a dimension, a unit of measure. These structures may depend on scale, e.g. on macroscopic scale condensed state of matter (and time) is well described as continuous medium, so we use dimensional units of length (and time). On the scale of atoms, the matter has a discrete structure, so we may count lattice sites and may not use a unit of length. If at small (e.g. at Plank) scale space (and/or time) is discrete, then we do not need a unit of length (time) for measuring, there is a fundamental length and we can just count.

Let us consider the following **discrete dynamics**:

$$S_{n+1} + S_{n-1} = \Phi(S_n), \tag{15}$$

which is obviously a (discrete) time (n) invertible in this implicit form. In the explicit form

$$S_{n+1} = F(S_n, S_{n-1}) = \Phi(S_n) - S_{n-1} \tag{16}$$

it is not obviously time invertible. If we take two step time lattice-make simplest discrete RD step and from one component-scalar S(n) construct two component-spinor $\Psi(n)$, we obtain explicit time invertible dynamics

$$\Psi_{n+1} = \Omega(\Psi_n), \ \Psi_{n+1} = \begin{pmatrix} S_{n+2} \\ S_{n+1} \end{pmatrix}, \ \Psi_n = \begin{pmatrix} S_n \\ S_{n-1} \end{pmatrix} \tag{17}$$

This **dynamical mechanism of origin spin** which connects time inversion symmetry and spin was invented when was constructed the theory of quanputers [Makhaldiani, 2011.2].

This mechanism indicates that with time inversion symmetry we can have only composed scalar fields. With the discovery of the Higgs particle with mass $125\ GeV$, a nice number $m_W/m_H \simeq 2/3$ appear, which, at least for me, indicates for composed nature of W and H, with a same mass of about $40\ GeV$ two and three valence constituents correspondingly. The fermion constituents ψ_n^a of W and scalar constituents φ_n^a of H compose scalar super multiplet (φ_n^a, ψ_n^a) with a flavor index n and color index a. Another notation is (h, sh)-(He, She:).

With exact SUSY we have cofinement by dimentional counting: superspace dimension is zero on the hadronic scale, hadrons are pointlike, color is confined inside hadrons. For SM QCD this picture indicates that at the hadronic scale we have effective SQCD, which contains scalar quarks.

The 40 GeV constituents may be good candidates in dark matter particles.
Coupling constant unification at $\alpha_u^{-1} = 29.0$ and scale $10^{16} GeV$ in MSSM [Makhaldani, 2014] has a relict on the SM scale: $\alpha_2^{-1}(m) = 29.0$ at $m = 41 GeV$.
If we extrapolate the SM value of $\alpha^{-1}(m_Z)$ to electron masse scale, we find $\alpha^{-1}(m_e) = 137.0$
Recent discovery of the second Higgs particle with mass $M_H = 750 GeV$ indicates an interesting structures. It is curious that $M_H/m_h = 750/125 = 6!$

In the Standard Model of Particle Physics (SM), the values of the coupling constants and masses of particles depends on the scale according to the Renormdynamic motion equations. One charge a, one mass m RD equations are

$$\begin{aligned} \dot{\alpha} &= \beta(\alpha), \\ \dot{m} &= \gamma(\alpha) m \end{aligned} \tag{18}$$

For the electron and nucleon masses, electrodynamic and pion-nucleon fine structure constants we have an empirical relation:

$$m_e/\alpha \simeq m_N/\alpha_{\pi N} \tag{19}$$

We take the relation $m/\alpha = const$ as an integral of renormdynamic motion equations for m and α, find exact form of the β function in the minimal mass parametrization

$$\begin{aligned} &\gamma(\alpha) = \gamma_1\alpha + \gamma_2\alpha^2 + ... = \gamma_1 A, \\ &A = f^{-1}(\alpha) = \alpha + \gamma_2/\gamma_1\alpha^2 + \gamma_3/\gamma_1\alpha^3 + ..., \\ &\alpha = f(A) = A + f_2A^2 + f_3A^3 + ... \end{aligned} \tag{20}$$

From the integral of motion, in the minimal mass parametrization: $\gamma(\alpha) = \gamma_1 \alpha$, we obtain

$$\begin{aligned}(\ln \alpha)^{\cdot} = (\ln m)^{\cdot} \Rightarrow \beta(\alpha)/\alpha = \gamma(\alpha) \\ = \gamma_1 \alpha \Rightarrow \beta(\alpha) = \beta_2 \alpha^2, \ \beta_2 = \gamma_1\end{aligned} \tag{21}$$

so, we have the following algebraic-diofant equations for the flavor and color content of the theory

$$\begin{aligned}\beta_n = 0, \ n \geq 3, \\ \beta_2 = \gamma_1\end{aligned} \tag{22}$$

and prediction for the dimension of space-time: $D = 4$. Solution of the motion equations are

$$\begin{aligned}\alpha(t) = \frac{\alpha_0}{1 - \alpha_0 \beta_2 t}, \\ m(t) = m_0 |\alpha_0^{-1} - \beta_2 t|^{-\gamma_1/\beta_2} = \frac{m_0}{\alpha_0} \alpha(t)\end{aligned} \tag{23}$$

In the multidimentional renormdynamics, when we have several (N) coupling constants and masses, we assume that there are maximal number $(N-1)$ integrals of motion H_n. If the number of integrals is N, we not have dynamics, we have only statics - finite field theory,
$\alpha_n = const,\ n = 1, ..., N.$
The idea of reduction to the one dimensional renormdynamics is simple:

$$\frac{d\alpha_n}{dt} = \beta_n(\alpha_1, ..., \alpha_{(N-1)}, \alpha_N) \Rightarrow \frac{d\alpha_n}{d\alpha} = B_n(\alpha_1, ..., \alpha_{(N-1)}, \alpha),\ \alpha = \alpha_N;$$
$$B_n(\alpha_1, ..., \alpha_{(N-1)}, \alpha) = \beta_n(\alpha_1, ..., \alpha_{(N-1)}, \alpha)/\beta_N(\alpha_1, ..., \alpha_{(N-1)}, \alpha),$$
$$\alpha_n = \sum_{k\geq 1} f_{nk}\alpha^k, \quad n = 1, 2, ..., N-1 \tag{24}$$

Koide Formula(s) (KF)

Koide's mass formula is an empirical relation among the charged lepton masses which holds with a striking precision,

$$Q = \frac{m_e + m_\mu + m_\tau}{(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2} \simeq \frac{2}{3} \tag{25}$$

were electron mass is $m_e = 0.510998910(13) MeV/c^2$, muon mass is $m_\mu = 105.658367(4) MeV/c^2$, tauon mass is $m_t = 1776.84(17) MeV/c^2$. The digits in parentheses are the uncertainties in the last figures. This gives $Q = 0.666659(10)$. Not only is this result odd in that three apparently random numbers should give a simple fraction, but also that Q is exactly halfway between the two extremes of $1/3$ (should the three masses be equal) and 1 (should one mass dominate). Similar matches have been found for quarks depending on running masses, and for triplets of quarks not of the same flavour. The Koide relation is exact (within experimental error) for the pole masses, which are low-energy quantities defined at different energy scales. It is RD integral of motion.

Under the generalized KF we mean the following formulas [Das, Makhaldiani 2016]

$$Q_n = \frac{m_1 + m_2 + ... + m_n}{(\sqrt{m_1} + \sqrt{m_2} + ... + \sqrt{m_n})^2} = q_n, \ q_3 \simeq \frac{2}{3} \tag{26}$$

In the solvable model of RD for coupling constants and masses, $m_k/\alpha_k = I = const$. So, the KF is equivalent to the corresponding relation for charges

$$Q_n = \frac{e_1^2 + e_2^2 + ... + e_n^2}{(e_1 + e_2 + ... + e_n)^2} = q_n, \ e_k = \sqrt{4\pi\alpha_k} \tag{27}$$

In the $n-$dimensional euklidian space a vector e with coordinates $(e_1, e_2, ..., e_n)$ has the length squire $e_1^2 + ... + e_n^2$. If all components are oriented in one dimension, the length is $(e_1 + ... + e_n)^2$.

If we take a symmetric vector $(1, 1, ..., 1)$ with length $\sqrt{n}$ than $\cos\varphi$ of the angler φ between charge vector $e = (e_1, e_2, ..., e_n)$ and the symmetric vector will be

$$\cos\varphi_n = \frac{e_1 + ... + e_n}{\sqrt{n}\sqrt{e_1^2 + ... + e_n^2}} = \frac{1}{\sqrt{nq_n}},$$
$$\cos\varphi_1 = \cos\varphi_2 = 1, \ \cos\varphi_3 = \frac{1}{\sqrt{2}} \tag{28}$$

If the sequence q_n is bounded, than $\cos\varphi_n \to 0$

Golden Ratio (RG)

RG can be defined as a solution of the following equation

$$x^2 + x - 1 = 0,$$
$$x = \frac{\sqrt{5}-1}{2} \tag{29}$$

The equation and the solution can be put in the following form

$$x = \frac{1}{1+x} = \frac{1}{1+\frac{1}{1+x}} = ... = \frac{1}{1+\frac{1}{1+...}} = \frac{\sqrt{5}-1}{2} = 0.618034 \tag{30}$$

We can consider corresponding recurrent relarion

$$x_{n+1} = \frac{1}{1+x_n},\ x_1 = 1,\ x_2 = 1/2,\ x_3 = 2/3,$$
$$x_4 = 3/5 = 0.6,\ x_5 = 5/8 = 0.625...$$
$$\lim_{n\to\infty} x_n = x = \frac{\sqrt{5}-1}{2} \tag{31}$$

So, the value of the Koide ratio corresponds to $x_3 = 2/3$. We might consider as an exact value of the ratio the GR. The recurrent relation is a discrete RD equation with the fixed point solution the GR.

The first higgs particle has a mass $m_h = 125GeV$, the second one has $m_H = 750GeV$. The two term Koide formula

$$Q_2 = \frac{m_h + m_H}{(\sqrt{m_h} + \sqrt{m_H})^2} = \frac{7}{(1+\sqrt{6})} = 0.588286 \lesssim \frac{2}{3} = 0.666667 \quad (32)$$

We suppose that there is a third higgs particle with mass $m_x = x$, which makes the Koide formula exact and find the mass:

$$Q_3 = \frac{m_h + m_H + m_x}{(\sqrt{m_h} + \sqrt{m_H} + \sqrt{m_x})^2} = \frac{2}{3},$$
$$3(7+x) = 2((a+\sqrt{x})^2) = 2a^2 + 4a\sqrt{x} + 2x \Downarrow$$
$$x - 4a\sqrt{x} - 2a^2 + 21 \Rightarrow \sqrt{x} = 2a \pm \sqrt{6a^2 - 21} = 13.9978 = 14,$$
$$a = 1 + \sqrt{6} = 3.44949,\ \ 2a = 6.89898,\ \ \sqrt{6a^2 - 21} = 7.09886$$
$$m_x = 196 m_h = 196 \times 125 = 24.5TeV. \quad (33)$$

If we take the second, negative, root $\sqrt{x}$,

$$\sqrt{x} = 2a - \sqrt{6a^2 - 21} = -0.199885,\ \ x = 0.039954,$$
$$m_x = 0.039954 m_h = 0.039954 \times 125 = 4.99425 = 5GeV. \quad (34)$$

Let us take Q_2 and find the second mass (charge)

$$\frac{1+x}{(1+\sqrt{x})^2} = \frac{2}{3} \Rightarrow x - 4\sqrt{x} + 1 = 0 \Rightarrow \sqrt{x} = 2 \pm \sqrt{3},$$
$$x_1 = (2+\sqrt{3})^2 = 7 + 4\sqrt{3} = 13.9282 \simeq 14, \; x_2 = (2-\sqrt{3})^2 = 7 - 4\sqrt{3}$$
$$m_1 = 125 \times 13.9282 = 1741.03 = 1741 GeV,$$
$$m_2 = 125 \times 0.0717968 = 8.9746 \simeq 9 GeV$$

Solitons are particlelike states, solutions of motion equations and they quantum extensions. Examples are solitons of SinGordon motion equation or barions-skirmions of Skyrme model. In particle theory, the skyrmion was described by Tony Skyrme in 1962 and consists of a quantum superposition of baryons and resonance states.
Skyrmions as topological objects are important in solid state physics. Researchers could read and write skyrmions using scanning tunneling microscopy. The topological charge, representing the existence and non-existence of skyrmions, can represent the bit states "1" and "0".

QCD consists of quarks and gluons. Quarks possess both color (r, g, b) and flavor $(u, d, s$,etc.), while gluons possess color (r, g, b) and anti-color $(\bar{r}, \bar{g}, \bar{b})$, but not flavor. An open **string** (a string with two endpoints) is ideally suited to account for such quantum numbers at its two ends. For quarks, one end represents color and the other end flavor. For gluons, one end represents color and the other anti-color. In string theory, there are **branes** (higher dimensional extended objects that are generalized membranes) to which the endpoints of an open string are confined. Applying this idea to QCD, we introduce N_c colored branes and N_f flavored branes at which open strings corresponding to quarks and gluons terminate. The energy of a string is given by the sum of the classical energy stored inside the string and the excitation energies of vibration and rotation. Because the classical energy of a string is proportional to its length and because gluons are massless, N_c colored branes should lie on top of one another. On the other hand, quarks possess intrinsic masses, and therefore the endpoints of a quark string, namely, a flavored brane and a colored brane should be separated from each other by a nonvanishing distance U. Then, the intrinsic quark mass m_q can be represented as $m_q = U \times$(string tension), where the string tension is the energy stored inside a unit length of string and is represented: string tension$=1/(2\pi\alpha')$ in terms of α' , historically called the Regge slope.

To describe QCD, we have to prepare Dp-branes and Dq-branes with p, q$\geq$3 for colored branes and flavored branes, respectively, and these branes should be located in the space of more than five dimensions. To evaluate the amplitude for a certain process to occur in the above picture, we have to sum up all the possible two-dimensional world sheets with the weight $\exp(iS)$, where the action S is given by S=(energy)$\times$(time)=(area of the strings world sheet)$/2\pi\alpha'$, following the Feymann path integral formulation.

Cumulative Effect: Production of particles from nuclei in a region, kinematically forbidden for reactions with free nucleons is connected to the existence of **Fluctons** - droplets of dense cold nuclear matter.
Classical fields have canonical, rational for integer D, (mass)dimensions e.g. in electrodynamics

$$L = \int d^D x(\bar{\psi}(\gamma(\partial - eA) - m)\psi - \frac{1}{4}F^2),$$
$$d_\psi = [\psi] = (D-1)/2,\ d_A = (D-2)/2,\ d_e = (4-D)/2 \quad (36)$$

Quantum corrections introduce (anomaly) corrections to the canonical dimensions, so the fields and coupling constants become **fractals.** At fixed points of RD, the fractals are self similar and their compositions present at low energies **unparticles.**

Qualitative picture of the (un)particle(like) objects we will illustrate with the simplest model of scalar field given by the following lagrangian

$$L = L(\Phi, M, \lambda, n) = \frac{1}{2}(\partial_\mu \Phi) - \frac{1}{2}M^2\Phi^2 - V(\Phi),\ \mu = 0, 1, 2, ..., D \quad (37)$$

where self interaction usually we take in the form

$$V(\Phi) = \lambda\Phi^n,\ n = -2, 1, 2, 3, 4, 6 \quad (38)$$

In renormalisible case,

$$n = \frac{2D}{D-2} = 2 + \varepsilon(D),\ \varepsilon(D) = \frac{4}{D-2},$$
$$D = \frac{2n}{n-2} = 2 + \varepsilon(n),\ \varepsilon(n) = \frac{4}{n-2}, \quad (39)$$

sometimes we consider also intermediate values of n and D and other forms of V.

In the free (self non interacting) field (particle) approximation: $\lambda = 0$, but in external gravitational field we have

$$L(g, \Phi, M) = \sqrt{-g} L(\Phi, M, 0), \ g = det g_{\mu\nu}(x) \tag{40}$$

Now we will see a nice composite particle mechanism :) Let us take a substitute: $\Phi = \varphi^k$, than we find

$$L(g, \Phi, M) = L((k\varphi^{k-1})^4 g, \varphi, M/k), \ g_{\mu\nu}(x) \Rightarrow (k\varphi^{k-1})^{4/D} g_{\mu\nu}(x) \tag{41}$$

Indeed

$$\begin{aligned} L(g, \Phi, M) &= \sqrt{-g}(k^2 \varphi^{2(k-1)} \frac{1}{2}(\partial_\mu \varphi)^2 - \frac{1}{2} M^2 \varphi^{2k}) \\ &= \sqrt{-g(k\varphi^{k-1})^4}(\frac{1}{2}(\partial_\mu \varphi)^2) - \frac{1}{2}(\frac{M}{k})^2 \varphi^2) \end{aligned} \tag{42}$$

Now, having an experience with constituent - composite particle relation, we turn attention on the self-interaction therm,

$$L = \sqrt{-g(k\varphi^{k-1})^4}(... - \frac{\lambda}{k^2}\varphi^N),\ N = kn - 2(k-1) \tag{43}$$

Most natural value of n for stable systems $(1+1 \to 1+1,\ 2 \to 2)$ is $n = 4$. In this case, $N = 2k+2$ and only natural value of constituents for which we have a renormalizable interaction is $k = 2 \Rightarrow N = 6$ with corresponding spacetime dimension $D = 3$. The most natural value for fission-fusion interaction $(1 \leftrightarrow 2)$ is $n = 3 \Rightarrow N = k+2$, for which we have realistic values $k = 2$ and $N = 4,\ D = 4$:) Other interesting values of naturally interpretable monomial (polynomial) interactions generally corresponds to the non-integer, fractional-fractal dimensions of space(time) D, with fractal-flucton-unparticle interpretations of the corresponding states of matter.

The size of particle-like states (solutions of the motion equations) are defined as $l \sim M^{-1}$, because at the boundary region, the linear part of the motion equations dominates and the Yukawa-like asymptotic $\Phi(r) \sim e^{-Mr}$ acts. In a pion-nucleon model for nucleon size we have $l_N \sim m_\pi^{-1} \simeq 1.43$ fm. The amplitude of the state (at maximum) $A \sim \lambda^{-\alpha},\ \alpha = 1/(n-2)$. Indeed, the motion equation do not contains the coupling constant after a scaling substitution $\Phi = \lambda^{-\alpha}\phi$, so a particle-like solution ϕ dos not contains λ and corresponding solution $\Phi = \lambda^{-\alpha}\phi \sim \lambda^{-\alpha}$,

$$\triangle\Phi + M^2\Phi + \lambda n\Phi^{n-1} = \lambda^{-\alpha}(\triangle\phi + M^2\phi + \lambda^{1-(n-2)\alpha} n\phi^{n-1}) = 0. \quad (44)$$

At not so low energies from string theory we may extract the following scalar field theory

$$L = \sqrt{-g}(\frac{1}{2}(\partial_\mu \Phi)^2 - \frac{1}{2}M^2\Phi^2 - \lambda\Phi^3),\ \mu = 0, 1, ..., D-1,\ D = 6 + \varepsilon \quad (45)$$

where $\varepsilon \in [0, 20]$. The one loop $\beta-$function is

$$\beta(a) = (D-6)a - \beta_2 a^2,\ a \sim \lambda^2 \quad (46)$$

and it has stable UV fixed point at $a = (D-6)/\beta_2$ and IR fixed point $a = 0$. Beyond this point we have an unparticle $\Phi = \phi^2$ with lagrangian

$$L = \sqrt{-g'}(\frac{1}{2}(\partial_\mu \phi)^2 - \frac{1}{2}(\frac{M}{2})^2\phi^2 - \frac{\lambda}{4}\phi^4),\ \mu = 0, 1, ..., D-1,$$
$$d = 4 - \varepsilon,\ \varepsilon \in [0, 1]. \quad (47)$$

The one loop $\beta-$function is

$$\beta(\lambda) = (d-4)\lambda + b\lambda^2 \tag{48}$$

and it has stable IR fixed point at $\lambda = (4-d)/b$. The UV fixed point is $\lambda = 0$. At this point we have reduction from higher dimentional Φ^3 to lower dimentional ϕ^4.
Another possibilities is an unparticle $\Phi = \varphi^4$ with lagrangian

$$L = \sqrt{-g''}(\frac{1}{2}(\partial_\mu\varphi)^2 - \frac{1}{2}(\frac{M}{4})^2\varphi^2 - \frac{\lambda}{16}\varphi^6),\ \mu = 0, 1, ..., d-1. \tag{49}$$

The one loop $\beta-$function is

$$\beta(\lambda) = (d-3)\lambda + c\lambda^2,\ d = 3-\varepsilon,\ \varepsilon \in [0, 2]. \tag{50}$$

The IR fixed point is $\lambda = \varepsilon/c$. UV fixed point is $\lambda = 0$.

Similar consideration gives reduction from higher energy ϕ^4 model to lower energy φ^6 one. Some technical questions remain. One of them concern to the substitution $\Phi = \phi^2$. It restricts Φ as $\Phi \geq 0$. OK, we already have a constraint, that the fields are real valued, we have a restriction

$$\phi^*(x) = \phi(x) = \frac{1}{(2\pi)^D} \int d^D p \exp(ipx)\hat{\phi}(p) \Rightarrow \hat{\phi}^*(p) = \hat{\phi}(-p) \qquad (51)$$

To formulate positivity condition is not so easy. We will take another path, we define the interaction as $\Phi^3 = (\Phi^2)^{3/2} \geq 0$. Then the substitution $\Phi^2 = \phi^4$ will works. Bytheway by this definition we made also another improvement: the potential become bounded from below. For the reduction the substitution $\Phi^2 = \phi^4$ also works,

$$L = \sqrt{-g}(\frac{1}{8\Phi^2})(\partial_\mu \Phi^2)^2 - \frac{1}{2}M^2\Phi^2 - \lambda(\Phi^2)^{n/2}, \; n = 3, 4. \qquad (52)$$

Note that by substitution

$$(\frac{\Phi}{\Phi_0})^2 = \phi^{2k},\ \phi^2 = \exp(\ln(\Phi^2/\Phi_0^2)/k)$$
$$= 1 + \frac{1}{k}\ln(\frac{\Phi}{\Phi_0})^2 + O(K^{-2}),\ \phi = \pm 1 + O(k^{-1}) \qquad (53)$$

we reduce the field theory to a discrete theory, to a system of bits. Also, changing dimension of space D and nonlinearity n restricted by condition

$$n = \frac{2D}{D-2},\ D = \frac{2n}{n-2},\ \frac{1}{n} + \frac{1}{D} = \frac{1}{2} \qquad (54)$$

we assume that they are functions of scale or coupling constant, due to monotonic property of the coupling constant. We have the following relation

$$\beta_n = -\frac{4}{(D-2)^2}\beta_D,$$
$$\beta_n = \mu\frac{dn}{d\mu} = \frac{dn}{d\lambda}\beta_\lambda,\ \beta_D = \mu\frac{dD}{d\mu} = \frac{dD}{d\lambda}\beta_\lambda. \qquad (55)$$

Let us consider a general dynamical system described by the following system of the ordinary differential equations [Arnold, 1978]

$$\dot{x}_n = v_n(x),\ 1 \leq n \leq N, \tag{56}$$

$\dot{x}_n$ stands for the total derivative with respect to the parameter t. When the number of the degrees of freedom is even, and

$$v_n(x) = \varepsilon_{nm}\frac{\partial H_0}{\partial x_m},\ 1 \leq n, m \leq 2M, \tag{57}$$

the system (56) is Hamiltonian one and can be put in the form

$$\dot{x}_n = \{x_n, H_0\}_0, \tag{58}$$

where the Poisson bracket is defined as

$$\{A, B\}_0 = \varepsilon_{nm}\frac{\partial A}{\partial x_n}\frac{\partial B}{\partial x_m} = A\frac{\overleftarrow{\partial}}{\partial x_n}\varepsilon_{nm}\frac{\overrightarrow{\partial}}{\partial x_m}B, \tag{59}$$

and summation rule under repeated indices has been used.

Let us consider the following Lagrangian

$$L = (\dot{x}_n - v_n(x))\psi_n \tag{60}$$

and the corresponding equations of motion

$$\dot{x}_n = v_n(x), \dot{\psi}_n = -\frac{\partial v_m}{\partial x_n}\psi_m. \tag{61}$$

The system (61) extends the general system (56) by linear equation for the variables ψ. The extended system can be put in the Hamiltonian form [Makhaldiani, Voskresenskaya, 1997]

$$\dot{x}_n = \{x_n, H_1\}_1, \dot{\psi}_n = \{\psi_n, H_1\}_1, \tag{62}$$

where first level (order) Hamiltonian is

$$H_1 = v_n(x)\psi_n \tag{63}$$

and (first level) bracket is defined as

$$\{A, B\}_1 = A(\frac{\overleftarrow{\partial}}{\partial x_n}\frac{\overrightarrow{\partial}}{\partial \psi_n} - \frac{\overleftarrow{\partial}}{\partial \psi_n}\frac{\overrightarrow{\partial}}{\partial x_n})B. \tag{64}$$

Note that when the Grassmann grading [Berezin, 1987] of the conjugated variables x_n and ψ_n are different, the bracket (64) is known as Buttin bracket[Buttin, 1996].
In the Faddeev-Jackiw formalism [Faddeev, Jackiw, 1988] for the Hamiltonian treatment of systems defined by first-order Lagrangians, i.e. by a Lagrangian of the form

$$L = f_n(x)\dot{x}_n - H(x), \tag{65}$$

motion equations

$$f_{mn}\dot{x}_n = \frac{\partial H}{\partial x_m}, \tag{66}$$

for the regular structure function f_{mn}, can be put in the explicit hamiltonian (Poisson; Dirac) form

$$\dot{x}_n = f^{-1}_{nm}\frac{\partial H}{\partial x_m} = \{x_n, x_m\}\frac{\partial H}{\partial x_m} = \{x_n, H\}, \tag{67}$$

where the fundamental Poisson (Dirac) bracket is

$$\{x_n, x_m\} = f^{-1}_{nm}, \;\; f_{mn} = \partial_m f_n - \partial_n f_m. \tag{68}$$

The system (61) is an important example of the first order regular hamiltonian systems. Indeed, in the new variables,

$$y_n^1 = x_n, y_n^2 = \psi_n, \tag{69}$$

lagrangian (60) takes the following first order form

$$\begin{aligned} L &= (\dot{x}_n - v_n(x))\psi_n \Rightarrow \frac{1}{2}(\dot{x}_n\psi_n - \dot{\psi}_n x_n) - v_n(x)\psi_n \\ &= \frac{1}{2}y_n^a \varepsilon^{ab}\dot{y}_n^b - H(y) \\ &= f_n^a(y)\dot{y}_n^a - H(y), f_n^a = \frac{1}{2}y_n^b\varepsilon^{ba}, H = v_n(y^1)y_n^2, \\ f_{nm}^{ab} &= \frac{\partial f_m^b}{\partial y_n^a} - \frac{\partial f_n^a}{\partial y_m^b} = \varepsilon^{ab}\delta_{nm}; \end{aligned} \tag{70}$$

corresponding motion equations and the fundamental Poisson bracket are

$$\dot{y}_n^a = \varepsilon_{ab}\delta_{nm}\frac{\partial H}{\partial y_m^b} = \{y_n^a, H\}, \{y_n^a, y_m^b\} = \varepsilon_{ab}\delta_{nm}. \tag{71}$$

To the canonical quantization of this system corresponds

$$[\hat{y}^a_n, \hat{y}^b_m] = i\hbar\varepsilon_{ab}\delta_{nm},\ \ \hat{y}^1_n = y^1_n,\ \ \hat{y}^2_n = -i\hbar\frac{\partial}{\partial y^1_n} \tag{72}$$

In this quantum theory, classical part, motion equations for y^1_n, remain classical.

Nambu dynamics

Nabu – Babylonian God
of Wisdom and Writing.

The Hamiltonian mechanics (HM) is in the fundamentals of mathematical description of the physical theories [Faddeev, Takhtajan, 1990]. But HM is in a sense blind; e.g., it does not make a difference between two opposites: the ergodic Hamiltonian systems (with just one integral of motion) [Sinai, 1993] and (super)integrable Hamiltonian systems (with maximal number of the integrals of motion).
Nabu mechanics (NM) [Nambu, 1973, Whittaker, 1927] is a proper generalization of the HM, which makes the difference between dynamical systems with different numbers of integrals of motion explicit (see, e.g.[Makhaldiani, 2007]).

Nambu dynamics

In the canonical formulation, the equations of motion of a physical system are defined via a Poisson bracket and a Hamiltonian, [Arnold, 1978]. In Nambu's formulation, the Poisson bracket is replaced by the Nambu bracket with $n+1, n \geq 1$, slots. For $n = 1$, we have the canonical formalism with one Hamiltonian. For $n \geq 2$, we have Nambu-Poisson formalism, with n Hamiltonians, [Nambu, 1973], [Whittaker, 1927].

The system of N vortexes can be described by the following system of differential equations, [Aref, 1983, Meleshko,Konstantinov, 1993]

$$\dot{z}_n = i \sum_{m \neq n}^{N} \frac{\gamma_m}{z_n^* - z_m^*}, \quad 1 \leq n \leq N, \tag{73}$$

where $z_n = x_n + iy_n$ are complex coordinate of the centre of n-th vortex, for $N = 3$, and the quantities

$$\begin{aligned} u_1 &= ln|z_2 - z_3|^2, \\ u_2 &= ln|z_3 - z_1|^2, \\ u_3 &= ln|z_1 - z_2|^2 \end{aligned} \tag{74}$$

reduce to the following system

$$\begin{aligned} \dot{u}_1 &= \gamma_1(e^{u_2} - e^{u_3}), \\ \dot{u}_2 &= \gamma_2(e^{u_3} - e^{u_1}), \\ \dot{u}_3 &= \gamma_3(e^{u_1} - e^{u_2}), \end{aligned} \tag{75}$$

The system (75) has two integrals of motion

$$H_1 = \sum_{i=1}^{3} \frac{e^{u_i}}{\gamma_i}, H_2 = \sum_{i=1}^{3} \frac{u_i}{\gamma_i}$$

and can be presented in the Nambu–Poisson form, [Makhaldiani, 1997,2]

$$\dot{u}_i = \omega_{ijk} \frac{\partial H_1}{\partial u_j} \frac{\partial H_2}{\partial u_k} = \{x_i, H_1, H_2\} = \omega_{ijk} \frac{e^{u_j}}{\gamma_j} \frac{1}{\gamma_k},$$

where

$$\omega_{ijk} = \epsilon_{ijk} \rho, \rho = \gamma_1 \gamma_2 \gamma_3$$

and the Nambu–Poisson bracket of the functions A, B, C on the three-dimensional phase space is

$$\{A, B, C\} = \omega_{ijk} \frac{\partial A}{\partial u_i} \frac{\partial B}{\partial u_j} \frac{\partial C}{\partial u_k}. \tag{76}$$

This system is superintegrable: for $N = 3$ degrees of freedom, we have maximal number of the integrals of motion $N - 1 = 2$.

The reduction of the dimensionless couplings in GUTs is achieved by searching for RD integrals of motion-renormdynamic invariant (RDI) relations among them holding beyond the unification scale. Finiteness results from the fact that there exist RDI relations among dimensional couplings that guarantee the vanishing of all beta-functions in certain GUTs even to all orders. In this case the number of the independent motion integrals N is equal to the number of the coupling constants. Note that in superintegrable dynamical systems the number of the integrals is $\leq N-1$, so the RD of the finite field theories is trivial, coupling constants do not run, the have fixed values, the renormdynamics is more than superintegrable, it is hyperintegrable. Developments in the soft supersymmetry breaking sector of GUTs and FUTs lead to exact RDI relations, i.e. reduction of couplings, in this dimensionful sector of the theory, too. Based on the above theoretical framework phenomenologically consistent FUTs have been constructed. The main goal expected from a unified description of interactions by the particle physics community is to understand the present day large number of free parameters of the SM in terms of a few fundamental ones. In other words, to achieve reduction of couplings at a more fundamental level.

As an example of the infinite dimensional Nambu-Poisson dynamics, let me conside the following extension of Schrödinger quantum mechanics [Makhaldiani, 2000]

$$iV_t = \Delta V - \frac{V^2}{2}, \tag{77}$$

$$i\psi_t = -\Delta\psi + V\psi. \tag{78}$$

An interesting solution to the equation for the potential (77) is

$$V = \frac{4(4-d)}{r^2}, \tag{79}$$

where d is the dimension of the spase. In the case of $d = 1$, we have the potential of conformal quantum mechanics.
The variational formulation of the extended quantum theory, is given by the following Lagrangian

$$L = (iV_t - \Delta V + \frac{1}{2}V^2)\psi. \tag{80}$$

The momentum variables are

$$P_v = \frac{\partial L}{\partial V_t} = i\psi, P_\psi = 0. \tag{81}$$

As Hamiltonians of the Nambu-theoretic formulation, we take the following integrals of motion

$$\begin{aligned} H_1 &= \int d^d x(\Delta V - \frac{1}{2}V^2)\psi, \\ H_2 &= \int d^d x(P_v - i\psi), \\ H_3 &= \int d^d x P_\psi. \end{aligned} \tag{82}$$

We invent unifying vector notation, $\phi = (\phi_1, \phi_2, \phi_3, \phi_4) = (\psi, P_\psi, V, P_v)$. Then it may be verified that the equations of the extended quantum theory can be put in the following Nambu-theoretic form

$$\phi_t(x) = \{\phi(x), H_1, H_2, H_3\}, \tag{83}$$

where the bracket is defined as

$$\{A_1, A_2, A_3, A_4\} = i\varepsilon_{ijkl} \int \frac{\delta A_1}{\delta\phi_i(y)} \frac{\delta A_2}{\delta\phi_j(y)} \frac{\delta A_3}{\delta\phi_k(y)} \frac{\delta A_4}{\delta\phi_l(y)} dy$$
$$= i \int \frac{\delta(A_1, A_2, A_3, A_4)}{\delta(\phi_1(y), \phi_2(y), \phi_3(y), \phi_4(y))} dy = i det(\frac{\delta A_k}{\delta\phi_l}). \qquad (84)$$

The basic building blocks of M theory are membranes and $M5-$branes. Membranes are fundamental objects carrying electric charges with respect to the 3-form C-field, and $M5$-branes are magnetic solitons. The Nambu-Poisson 3-algebras appear as gauge symmetries of superconformal Chern-Simons nonabelian theories in $2+1$ dimensions with the maximum allowed number of $N=8$ linear supersymmetries.
The Bagger and Lambert [Bagger, Lambert, 2007] and, Gustavsson [Gustavsson, 2007] (BLG) model is based on a 3-algebra,

$$[T^a, T^b, T^c] = f_d^{abc} T^d \tag{85}$$

where T^a, are generators and f_{abcd} is a fully anti-symmetric tensor.

Given this algebra, a maximally supersymmetric Chern-Simons lagrangian is:

$$L = L_{CS} + L_{matter},$$
$$L_{CS} = \frac{1}{2}\varepsilon^{\mu\nu\lambda}(f_{abcd}A^{ab}_{\mu}\partial_{\nu}A^{cd}_{\lambda} + \frac{2}{3}f_{cdag}f^{g}_{efb}A^{ab}_{\mu}A^{cd}_{\nu}A^{ef}_{\lambda}), \quad (86)$$

$$L_{matter} = \frac{1}{2}B^{Ia}_{\mu}B^{\mu I}_{a} - B^{Ia}_{\mu}D^{\mu}X^{I}_{a}$$
$$+\frac{i}{2}\bar{\psi}^{a}\Gamma^{\mu}D_{\mu}\psi_{a} + \frac{i}{4}\bar{\psi}^{b}\Gamma_{IJ}x^{I}_{c}x^{J}_{d}\psi_{a}f^{abcd}$$
$$-\frac{1}{12}tr([X^{I},X^{J},X^{K}][X^{I},X^{J},X^{K}]),\ I = 1,2,...,8, \quad (87)$$

where A^{ab}_{μ} is gauge boson, ψ^{a} and $X^{I} = X^{I}_{a}T^{a}$ matter fields. If $a = 1,2,3,4$, then we can obtain an $SO(4)$ gauge symmetry by choosing $f_{abcd} = f\varepsilon_{abcd}$, f being a constant. It turns out to be the only case that gives a gauge theory with manifest unitarity and $N = 8$ supersymmetry.

The action has the first order form so we can use the formalism of the first section. The motion equations for the gauge fields

$$f^{nm}_{abcd}\dot{A}^{cd}_{m}(t,x)=\frac{\delta H}{\delta A^{ab}_{n}(t,x)},\ f^{nm}_{abcd}=\varepsilon^{nm}f_{abcd} \tag{88}$$

take canonical form

$$\dot{A}^{ab}_{n}=f^{abcd}_{nm}\frac{\delta H}{\delta A^{cd}_{m}}=\{A^{ab}_{n},A^{cd}_{m}\}\frac{\delta H}{\delta A^{cd}_{m}}=\{A^{ab}_{n},H\},$$
$$\{A^{ab}_{n}(t,x),A^{cd}_{m}(t,y)\}=\varepsilon_{nm}f^{abcd}\delta^{(2)}(x-y) \tag{89}$$

The quasi-classical description of the motion of a relativistic (nonradiating) point particle with spin in accelerators and storage rings includes the equations of orbit motion

$$\dot{x}_n = f_n(x),\ \ f_n(x) = \varepsilon_{nm}\partial_m H,\ \ n,m = 1,2,...,6;$$
$$x_n = q_n,\ \ x_{n+3} = p_n,\ \ \varepsilon_{n,n+3} = 1,\ \ n = 1,2,3;$$
$$H = e\Phi + c\sqrt{\wp^2 + m^2c^2},\ \ \wp_n = p_n - \frac{e}{c}A_n \qquad (90)$$

and Thomas-BMT equations
[Tomas, 1927, Bargmann, Michel,Telegdi, 1959] of classical spin motion

$$\dot{s}_n = \varepsilon_{nmk}\Omega_m s_k = \{H_1, H_2, s_n\},\ \ H_1 = \Omega\cdot s,\ \ H_2 = s^2,$$
$$\{A,B,C\} = \varepsilon_{nmk}\partial_n A\partial_m B\partial_k C, \qquad (91)$$

$$\Omega_n = \frac{-e}{m\gamma c}((1+k\gamma)B_n - k\frac{(B\cdot\wp)\wp_n}{m^2c^2(1+\gamma)} + \frac{1+k(1+\gamma)}{mc(1+\gamma)}\varepsilon_{nmk}E_m\wp_k) \tag{92}$$

where, parameters e and m are the charge and the rest mass of the particle, c is the velocity of light, $k=(g-2)/2$ quantifies the anomalous spin g factor, γ is the Lorentz factor, p_n are components of the kinetic momentum vector, E_n and B_n are the electric and magnetic fields, and A_n and Φ are the vector and scalar potentials;

$$B_n = \varepsilon_{nmk}\partial_m A_k,\ E_n = -\partial_n\Phi - \frac{1}{c}\dot{A}_n,$$
$$\gamma = \frac{H-e\Phi}{mc^2} = \sqrt{1+\frac{\wp^2}{m^2c^2}} \tag{93}$$

The spin motion equations we put in the Nambu-Poisson form. Hamiltonization of this dynamical system according to the general approach of the previous sections we will put in the ground of the optimal control theory of the accelerator.

The general method of Hamiltonization of the dynamical systems we can use also in the spinning particle case. Let us invent unified configuration space $q = (x, p, s),\ x_n = q_n,\ p_n = q_{n+3},\ s_n = q_{n+6},\ n = 1, 2, 3$; extended phase space, (q_n, ψ_n) and hamiltonian

$$H = H(q, \psi) = v_n \psi_n,\ n = 1, 2, ...9; \tag{94}$$

motion equations

$$\begin{aligned} \dot{q}_n &= v_n(q), \\ \dot{\psi}_n &= -\frac{\partial v_m}{\partial q_n}\psi_m \end{aligned} \tag{95}$$

where the velocities v_n depends on external fields as in previous section as control parameters which can be determined according to the optimal control criterium.

Electric Dipole Moments (EDM) of Protons and Deuterons

EDM are one of the keys to understand the origin of our Universe [Sakharov, 1967]. Andrei Sakharov formulated three conditions for baryogenesis:
1. Early in the evolution of the universe, the baryon number conservation must be violated sufficiently strongly,
2. The C and CP invariances, and T invariance thereof, must be violated, and
3. At the moment when the baryon number is generated, the evolution of the universe must be out of thermal equilibrium.
CP violation in kaon decays is known since 1964, it has been observed in B-decays and charmed meson decays. The Standard Model (SM) accommodates CP violation via the phase in the Cabibbo-Kobayashi-Maskawa matrix.
CP and P violation entail nonvanishing P and T violating electric dipole moments (EDM) of elementary particles $\vec{d} = d\vec{s}$.

Electric Dipole Moments (EDM) of Protons and Deuterons

Although extremely successful in many aspects, the SM has at least two weaknesses: neutrino oscillations do require extensions of the SM and, most importantly, the SM mechanisms fail miserably in the expected baryogenesis rate.
Simultaneously, the SM predicts an exceedingly small electric dipole moment of nucleons $10^{-33} < d_n < 10^{-31} e \cdot cm$, way below the current upper bound for the neutron EDM, $d_n < 2.9 \times 10^{-26} e \cdot cm$. In the quest for physics beyond the SM one could follow either the high energy trail or look into new methods which offer very high precision and sensitivity.
Supersymmetry is one of the most attractive extensions of the SM and S. Weinberg emphasized [Weinberg, 1993]: "Endemic in supersymmetric (SUSY) theories are CP violations that go beyond the SM. For this reason it may be that the next exciting thing to come along will be the discovery of a neutron electric dipole moment."

Electric Dipole Moments (EDM) of Protons and Deuterons

The SUSY predictions span typically $10^{-29} < d_n < 10^{-24} e \cdot cm$ and precisely this range is targeted in the new generation of EDM searches [Roberts, Marciano, 2010]. There is consensus among theorists that measuring the EDM of the proton, deuteron and helion is as important as that of the neutron. Furthermore, it has been argued that T-violating nuclear forces could substantially enhance nuclear EDM [Flambaum, Khriplovich, Sushkov, 1986]. At the moment, there are no significant direct upper bounds available on d_p or d_d. Non-vanishing EDMs give rise to the precession of the spin of a particle in an electric field. In the rest frame of a particle

$$\dot{s}_n = \varepsilon_{nmk}(\Omega_m s_k + d_m E_k), \ \Omega_m = -\mu B_m, \tag{96}$$

where in terms of the lab frame fields

$$\begin{aligned} B_n &= \gamma(B^l_n - \varepsilon_{nmk}\beta_m E^l_k), \\ E_n &= \gamma(E^l_n + \varepsilon_{nmk}\beta_m B^l_k) \end{aligned} \tag{97}$$

Now we can apply the Hamiltonization and optimal control theory methods to this dynamical system.

Note that the procedure of reduction of the higher order dynamical system, e.g. second order Euler-Lagrange motion equations, to the first order dynamical systems, in the case to the Hamiltonian motion equations, can be continued using fractal calculus. E.g. first order system can be reduced to the half order one,

$$\begin{aligned} &D^{1/2}q = \psi, \\ &D^{1/2}\psi = p \Leftrightarrow \dot{q} = p. \end{aligned} \tag{98}$$

Discrete dynamical systems and Quanputers

Computers are physical devices and their behavior is determined by physical laws. The Quantum Computations [Benenti, Casati, Strini, 2004 , Nielsen, Chuang, 2000], Quantum Computing, Quanputing [Makhaldiani, 2007.2], is a new interdisciplinary field of research, which benefits from the contributions of physicists, computer scientists, mathematicians, chemists and engineers. Contemporary digital computer and its logical elements can be considered as a spatial type of discrete dynamical systems [Makhaldiani, 2001]

$$S_n(k+1) = \Phi_n(S(k)), \tag{99}$$

where

$$S_n(k), \quad 1 \leq n \leq N(k), \tag{100}$$

is the state vector of the system at the discrete time step k. Vector S may describe the state and Φ transition rule of some Cellular Automata [Toffoli, Margolus, 1987].The systems of the type (99) appears in applied mathematics as an explicit finite difference scheme approximation of the equations of the physics [Samarskii, Gulin, 1989].

Definition: *We assume that the system (99) is time-reversible if we can define the reverse dynamical system*

$$S_n(k) = \Phi_n^{-1}(S(k+1)). \tag{101}$$

In this case the following matrix

$$M_{nm} = \frac{\partial \Phi_n(S(k))}{\partial S_m(k)}, \tag{102}$$

is regular, i.e. has an inverse. If the matrix is not regular, this is the case, for example, when $N(k+1) \neq N(k)$, we have an irreversible dynamical system (usual digital computers and/or corresponding irreversible gates).

Let us consider an extension of the dynamical system (99) given by the following action function

$$A = \sum_{kn} l_n(k)(S_n(k+1) - \Phi_n(S(k))) \tag{103}$$

and corresponding motion equations

$$\begin{aligned} S_n(k+1) &= \Phi_n(S(k)) = \frac{\partial H}{\partial l_n(k)}, \\ l_n(k-1) &= l_m(k)\frac{\partial \Phi_m(S(k))}{\partial S_n(k)} = l_m(k)M_{mn}(S(k)) = \frac{\partial H}{\partial S_n(k)}, \end{aligned} \tag{104}$$

where

$$H = \sum_{kn} l_n(k)\Phi_n(S(k)), \tag{105}$$

is discrete Hamiltonian. In the regular case, we put the system (104) in an explicit form

$$\begin{aligned} S_n(k+1) &= \Phi_n(S(k)), \\ l_n(k+1) &= l_m(k)M^{-1}_{mn}(S(k+1)). \end{aligned} \tag{106}$$

From this system it is obvious that, when the initial value $l_n(k_0)$ is given, the evolution of the vector $l(k)$ is defined by evolution of the state vector $S(k)$. The equation of motion for $l_n(k)$ - Elenka is linear and has an important property that a linear superpositions of the solutions are also solutions.

Statement: *Any time-reversible dynamical system (e.g. a time-reversible computer) can be extended by corresponding linear dynamical system (quantum - like processor) which is controlled by the dynamical system and has a huge computational power,*
[Makhaldiani, 2001, Makhaldiani, 2002, Makhaldiani, 2007.2, Makhaldiani, 2011.2].

(de)Coherence criterion

For motion equations (104) in the continual approximation, we have

$$
\begin{aligned}
&S_n(k+1) = x_n(t_k+\tau) = x_n(t_k) + \dot{x}_n(t_k)\tau + O(\tau^2),\\
&\dot{x}_n(t_k) = v_n(x(t_k)) + O(\tau), \quad t_k = k\tau,\\
&v_n(x(t_k)) = (\Phi_n(x(t_k)) - x_n(t_k))/\tau;\\
&M_{mn}(x(t_k)) = \delta_{mn} + \tau\frac{\partial v_m(x(t_k))}{\partial x_n(t_k)}.
\end{aligned}
\tag{107}
$$

(de)Coherence criterion: *the system is reversible, the linear (quantum, coherent, soul) subsystem exists, when the matrix M is regular,*

$$
detM = 1 + \tau\sum_n \frac{\partial v_n}{\partial x_n} + O(\tau^2) \neq 0. \tag{108}
$$

For the Nambu - Poisson dynamical systems (see e.g. [Makhaldiani, 2007])

$$
\begin{aligned}
&v_n(x) = \varepsilon_{nm_1m_2\ldots m_p}\frac{\partial H_1}{\partial x_{m_1}}\frac{\partial H_2}{\partial x_{m_2}}\ldots\frac{\partial H_p}{\partial x_{m_p}}, \quad p = 1,2,3,\ldots,N-1,\\
&\sum_n \frac{\partial v_n}{\partial x_n} \equiv divv = 0.
\end{aligned}
\tag{109}
$$

Construction of the reversible discrete dynamical systems

Let me motivate an idea of construction of the reversible dynamical systems by simple example from field theory. There are renormalizable models of scalar field theory of the form (see, e.g. [Makhaldiani, 1980])

$$L = \frac{1}{2}(\partial_\mu \varphi \partial^\mu \varphi - m^2 \varphi^2) - g\varphi^n, \tag{110}$$

with the constraint

$$n = \frac{2d}{d-2}\ , \tag{111}$$

where d is dimension of the space-time and n is degree of nonlinearity. It is interesting that if we define d as a function of n, we find

$$d = \frac{2n}{n-2} \tag{112}$$

the same function !
Thing is that, the constraint can be put in the symmetric implicit form [Makhaldiani, 1980]

$$\frac{1}{n} + \frac{1}{d} = \frac{1}{2} \tag{113}$$

Generalization of the idea

Now it is natural to consider the following symmetric function

$$f(y) + f(x) = c \tag{114}$$

and define its solution

$$y = f^{-1}(c - f(x)). \tag{115}$$

This is the general method, that we will use in the following construction of the reversible dynamical systems. In the simplest case,

$$f(x) = x, \tag{116}$$

we take

$$y = S(k+1), \quad x = S(k-1), \quad c = \tilde{\Phi}(S(k)) \tag{117}$$

and define our reversible dynamical system from the following symmetric, implicit form (see also [Toffoli, Margolus, 1987])

$$S(k+1) + S(k-1) = \tilde{\Phi}(S(k)), \tag{118}$$

explicit form of which is

$$\begin{aligned} S(k+1) &= \Phi(S(k), S(k-1)) \\ &= \tilde{\Phi}(S(k)) - S(k-1). \end{aligned} \tag{119}$$

Generalization of the idea

This dynamical system defines given state vector by previous two state vectors. We have reversible dynamical system on the time lattice with time steps of two units,

$$\begin{aligned} &S(k+2,2) = \Phi(S(k,2)),\\ &S(k+2,2) \equiv (S(k+2), S(k+1)),\\ &S(k,2) \equiv (S(k), S(k-1))). \end{aligned} \tag{120}$$

Internal, spin, degrees of freedom

Starting from a general discrete dynamical system, we obtained reversible dynamical system with internal(spin,bit) degrees of freedom

$$\begin{aligned} S_{ns}(k+2) &\equiv \begin{pmatrix} S_n(k+2) \\ S_n(k+1) \end{pmatrix} = \begin{pmatrix} \Phi_n(\Phi(S(k)) - S(k-1)) - S(k)) \\ \Phi_n(S(k)) - S_n(k-1) \end{pmatrix} \\ &\equiv \Phi_{ns}(S(k)), \quad s = 1, 2 \end{aligned} \tag{121}$$

where

$$S(k) \equiv (S_{ns}(k)), \quad S_{n1}(k) \equiv S_n(k), \quad S_{n2}(k) \equiv S_n(k-1) \tag{122}$$

For the extended system we have the following action

$$A = \sum_{kns} l_{ns}(k)(S_{ns}(k+2) - \Phi_{ns}(S(k))) \tag{123}$$

and corresponding motion equations

$$S_{ns}(k+2) = \Phi_{ns}(S(k)) = \frac{\partial H}{\partial l_{ns}(k)},$$
$$l_{ns}(k-2) = l_{mt}(k)\frac{\partial \Phi_{mt}(S(k))}{\partial S_{ns}(k)}$$
$$= l_{mt}(k)M_{mtns}(S(k)) = \frac{\partial H}{\partial S_{ns}(k)}, \quad (124)$$

By construction, we have the following reversible dynamical system

$$S_{ns}(k+2) = \Phi_{ns}(S(k)),$$
$$l_{ns}(k+2) = l_{mt}(k)M^{-1}_{mtns}(S(k+2)), \quad (125)$$

with classical S_{ns} and quantum l_{ns}(in the external, background S) string bit dynamics.

p-point cluster and higher spin states reversible dynamics, or pit string dynamics

We can also consider p-point generalization of the previous structure,

$$f_p(S(k+p)) + f_{p-1}(S(k+p-1)) + ... + f_1(S(k+1))$$
$$+f_1(S(k-1)) + ... + f_p(S(k-p)) = \tilde{\Phi}(S(k)),$$
$$S(k+p) = \Phi(S(k), S(k+p-1), ..., S(k-p))$$
$$\equiv f_p^{-1}(\tilde{\Phi}(S(k)) - f_{p-1}(S(k+p-1)) - ... - f_p(S(k-p))) \quad (126)$$

and corresponding reversible p-oint cluster dynamical system

$$S(k+p,p) \equiv \Phi(S(k,p)),$$
$$S(k+p,p) \equiv (S(k+p), S(k+p-1), ..., S(k+1)),$$
$$S(k,p) \equiv (S(k), S(k-1), ..., S(k-p+1)), \quad S(k,1) = S(k). \quad (127)$$

So we have general method of construction of the reversible dynamical systems on the time (tame) scale p. The method of linear extension of the reversible dynamical systems (see [Makhaldiani, 2001] and previous section) defines corresponding Quanputers,

$$S_{ns}(k+p) = \Phi_{ns}(S(k)),$$
$$l_{ns}(k+p) = l_{mt}(k) M^{-1}_{mtns}(S(k+p)), \quad (128)$$

p-point cluster and higher spin states reversible dynamics, or pit string dynamics

This case the quantum state function $l_{ns},\ s = 1, 2, ...p$ will describes the state with spin $(p-1)/2$.
Note that, in this formalism for reversible dynamics minimal value of the spin is $1/2$. There is not a place for a scalar dynamics, or the scalar dynamics is not reversible. In the Standard model (SM) of particle physics, [Beringer et al, 2012], all of the fundamental particles, leptons, quarks and gauge bosons have spin. Only scalar particles of the SM are the Higgs bosons. Perhaps the scalar particles are composed systems or quasiparticles like phonon, or Higgs dynamics is not reversible (a mechanism for 'time arrow').

Social profit of big collaborations

Nowadays there are several big collaborations in science, e.g. LHC. Scientific value of LHC depends on three components, the highest quality of accelerator, highest quality of detectors and distributed data processing. The first two components need good mathematical and physical modeling. Third component and the collaboration as a social structure are not under (anther) the control by scientific methods and corresponding modeling. By definition, scientific collaborations (SC) have a main scientific aim: to obtain answer on the important scientific question(s) and maybe gain extra scientific bonus: new important questions and discoveries. SC is more open information system than e.g. finance or military systems. So, it is possible to describe and optimize SC by scientific methods. Profit from scientific modeling of SC maybe also for other information systems and social structures.

GRID and Quanputing

As an example of GRID we take LHC Computing Grid. The LHC Computing Grid (LCG), is an international collaborative project that consists of a grid-based computer network infrastructure incorporating over 170 computing centers in 36 countries. It was designed by CERN to handle the prodigious volume of data produced by Large Hadron Collider (LHC) experiments. The Large Hadron Collider at CERN was designed to prove or disprove the existence of the Higgs boson, an important but elusive piece of knowledge that had been sought by particle physicists for over 40 years. A very powerful particle accelerator was needed, because Higgs bosons might not be seen in lower energy experiments, and because vast numbers of collisions would need to be studied. Such a collider would also produce unprecedented quantities of collision data requiring analysis. Therefore, advanced computing facilities were needed to process the data. A design report was published in 2005. It was announced to be ready for data on 3 October 2008. It incorporates both private fiber optic cable links and existing high-speed portions of the public Internet. At the end of 2010, the Grid consisted of some 200,000 processing cores and 150 petabytes of disk space, distributed across 34 countries.

The data stream from the detectors provides approximately 300 GByte/s of data, which after filtering for "interesting events", results in a data stream of about 300 MByte/s. The CERN computer center, considered "Tier 0" of the LHC Computing Grid, has a dedicated 10 Gbit/s connection to the counting room. The project was expected to generate 27 TB of raw data per day, plus 10 TB of "event summary data", which represents the output of calculations done by the CPU farm at the CERN data center. This data is sent out from CERN to eleven Tier 1 academic institutions in Europe, Asia, and North America, via dedicated 10 Gbit/s links. This is called the LHC Optical Private Network. More than 150 Tier 2 institutions are connected to the Tier 1 institutions by general-purpose national research and education networks. The data produced by the LHC on all of its distributed computing grid is expected to add up to 10-15 PB of data each year. In total, the four main detectors at the LHC produced 13 petabytes of data in 2010. The Tier 1 institutions receive specific subsets of the raw data, for which they serve as a backup repository for CERN. They also perform reprocessing when recalibration is necessary. In 2015, CERN switched away from Scientific Linux to CentOS. Distributed computing resources for analysis by end-user physicists are provided by the Open Science Grid, Enabling Grids for E-sciencE, and LHC@home projects, http://wlcg.web.cern.ch/. Update of the Computing Models of the WLCG and the LHC Experiments:
http://cds.cern.ch/record/1695401/files/LCG-TDR-002.pdf

The idea of computations on quanputers is in finding of the needed (value of the) state (wave function $\psi(t,x)$) from the initial, easy constructible, state ($\psi(0,x),$) which is superposition of different states, including interesting one, with the same weight. During the computation the weight of the interesting state is growing till the value when we can guess the solution of the problem and then test it, which is much more easier then to find it.
Let us consider the following nonlinear evolution equation

$$iV_t = \Delta V - \frac{1}{2}V^2 + J, \tag{129}$$

extended Lagrangian and Hamiltonian

$$L = \int dx^D (iV_t - \Delta V + \frac{1}{2}V^2 - J)\psi,$$
$$H = \int dx^D (\Delta V - \frac{1}{2}V^2 + J)\psi \tag{130}$$

and corresponding Hamiltonian motion equations

$$\begin{aligned} &iV_t = \Delta V - \frac{1}{2}V^2 + J = \{V, H\}, \\ &i\psi_t = -\Delta\psi + V\psi = \{\psi, H\}, \\ &\{V(t,x), \psi(t,y)\} = \delta^D(x-y) \end{aligned} \tag{131}$$

The solution of the problem is given in the form

$$|T) = U(T)|0), \;\; \psi(t,x) =< x|t), \;\; U(T) = Pexp(-i\int_0^T dtH(t)) \tag{132}$$

Under the programming of the quanputer we understand construction of the potential V, or the corresponding Hamiltonian. For the given potential, we calculate corresponding source J.
The discrete version of the system can be put in the form

$$\begin{aligned} &S_m(n+1) = \Phi_n(S(n)) + J_m(n), \\ &\Psi_m(n-1) = A_{mk}(S(n))\Psi_k(n), \;\; A_{mk}(S(n)) = \frac{\partial\Phi_k(S(n))}{\partial S_m(n)} \end{aligned} \tag{133}$$

or, in the regular case, when the matrix A is regular,

we obtain explicit form of the corresponding discrete dynamics

$$S_m(n+1) = \Phi_n(S(n)) + J_m(n),$$
$$\Psi_m(n+1) = A^{-1}_{mk}(S(n+1))\Psi_k(n), \tag{134}$$

Now the state vector $S(n)$ and wave vector $\Psi_m(n)$ may correspond not only to the discrete values of the potential $V(n,m) = S_m(n)$, and wave function $\psi(n,m) = \Psi_m(n)$ but also any representation of the computing process from theoretical to experimental realization on a quanputer, including algorithm of solution, higher level programm realization of the algorithm. **Today, without big efforts, we can modify (some) GRID elements in time-invertible form. After development of the quanputer technologies, we can modify (some) GRID elements in quanputer forms.**

A way to the Solution of the Traveling salesman problem (TSP) with Quanputing

The $NP \stackrel{?}{=} P$ problem will be solved if for some $NP-$ complete problem, e.g. TSP, a polynomial algorithm find; or show that there is not such an algorithm; or show that it is impossible to find definite answer to that question.
TSP means to find minimal length path between N fixed points on a surface, which attends any point ones. We consider a system where N points with quenched positions $x_1, x_2, ..., x_N$ are independently distributed on a finite domain D with a probability density function $p(x)$. In general, the domain D is multidimensional and the points x_n are vectors in the corresponding Euclidean space. Inside the domain D we consider a polymer chain composed of N monomers whose positions are denoted by $y_1, y_2, ..., y_N$. Each monomer y_n is attached to one of the quenched sites x_m and only one monomer can be attached to each site. The state of the polymer is described by a permutation $\sigma \in \Sigma_N$ where Σ_N is the group of permutations of N objecs.

The Hamiltonian for the system is given by

$$H = \sum_{n=1}^{N} V(|y_n - y_{n-1}|) \tag{135}$$

Here V is the interaction between neighboring monomers on the polymer chain. For convenience the chain is taken to be closed, thus we take the periodic boundary condition $x_0 = x_N$. A physical realization of this system is one where the x_n are impurities where the monomers of a polymer loop are pinned. In combinatorial optimization, if one takes $V(x)$ to be the norm, or distance, of the vector x then $H(\sigma)$ is the total distance covered by a path which visits each site x_n exactly once. The problem of finding σ_0 which minimizes $H(\sigma)$ is known as the traveling salesman problem (TSP) [Gutin, Pannen, 2002].

In field theory language to the TSP we correspond the calculation of the following correlator

$$G_{2N}(x_1,x_2,...,x_N) = Z_0^{-1}\int d\varphi(x)\varphi^2(x_1)\varphi^2(x_2)...\varphi^2(x_N)e^{-S(\varphi)}$$
$$= \frac{\delta^{2N}F(J)}{\delta J(x_1)^2...\delta J(x_N)^2},\ F(J)=\ln Z(J),$$
$$Z(J) = \int d\varphi e^{-\frac{1}{2}\varphi\cdot A\cdot\varphi + J\cdot\varphi} = e^{\frac{1}{2}J\cdot A^{-1}\cdot J},\ A^{-1}(x,y;m) = e^{-m|x-y|},$$
$$L_{min}(x_1,...,x_N) = -\frac{d}{dm}\ln G_{2Ns} + O(e^{-am})$$
$$< A^{-1} > \equiv \frac{1}{\Gamma(s)}\int_0^\infty dm m^{s-1}A^{-1}(x,y;m) = \frac{1}{|x-y|^s}$$
$$= L_s A^{-1}(x-y;s)$$
$$k(d)\Delta_d L_s A^{-1}(x;s) = \delta^d(x) \Rightarrow A(x;s) = k(d)\Delta_d L_s,$$
$$s = d-2; \varphi = \varphi(x,m). \qquad (136)$$

If we take relativistic massive scalar field, then $A = \Delta_d + m^2$,

$$A^{-1}(x) \sim |x|^{2-d} e^{-m|x|}, \tag{137}$$

and for $d = 2$, we also have the needed behaviour. Note that G_{2N} is symmetric with respect to its arguments and contains any paths including minimal length one.

We consider a novel method to generate a polynomial expression for each of the Euler sums,

$$E_k = \sum_{n=1}^{N} n^k, \ k \in Z^+ (k = 0, 1, 2, ...) \tag{138}$$

One of the way of calculation of the sum

$$E_k(N) = \frac{N^{k+1}}{k+1} + P_k(N), \ P_k = x_k N^k + x_{k-1} N^{k-1} + ... + x_0 \tag{139}$$

we show by explicit calculation of E_2.
For particular values $N = 1, \ 2$ and 3, we have

$$\begin{aligned} x_2 + x_1 + x_0 &= 1 - 1/3 = 2/3, \\ 4x_2 + 2x_1 + x_0 &= 5 - 8/3 = 7/3, \\ 9x_2 + 3x_1 + x_0 &= 14 - 27/3 = 5 \end{aligned} \tag{140}$$

Subtracting from the second equation the first and from third the second, we obtain

$$3x_2 + x_1 = 5/3,$$
$$5x_2 + x_1 = 8/3 \tag{141}$$

than we have

$$2x_2 = 1 \Rightarrow x_2 = 1/2 \Downarrow$$
$$x_1 = 5/3 - 3x_2 = 1/6 \Rightarrow x_0 = 2/3 - x_1 - x_2 = 2/3 - 1/6 - 1/2 = 0, \Downarrow$$
$$E_3(N) = N^3/3 + N^2/2 + N/6$$
$$= N(N+1)(2N+1)/6 = N(N+1/2)(N+1)/3 \tag{142}$$

Note that, the right hand side have a sense also for $N \leq 0$ and has zeros at $N = 0, -1/2, -2$.

For general case $E_k(N)$ we have

$$x_k(N) = detV_l(N)/detW_k(N),$$

$$detW_k(N) = det \begin{pmatrix} 1 & N_1 & . & N_1^k \\ 1 & N_2 & . & N_2^k \\ . & . & . & . \\ 1 & N_{k+1} & . & N_{k+1}^k \end{pmatrix},$$

$$detV_l(N) = det \begin{pmatrix} 1 & N_1 & . & N_1^k \\ 1 & N_2 & . & N_2^k \\ \bar{E}_l(N_1) & \bar{E}_l(N_2) & . & \bar{E}_l(N_{k+1}) \\ 1 & N_{k+1} & . & N_{k+1}^k \end{pmatrix}, \; X = \begin{pmatrix} x_0 \\ x_1 \\ . \\ x_k \end{pmatrix}$$

$$\bar{E}_k(N_l) = E_k(N_l) - \frac{N_l^{k+1}}{k+1}, \; E = \begin{pmatrix} \bar{E}_k(N_1) \\ \bar{E}_k(N_2) \\ . \\ \bar{E}_k(N_{k+1}) \end{pmatrix},$$

$$WX = E, \; X = W^{-1}E \tag{143}$$

As a numbers N_n we can take any different integers, but the simplest choice is: $N_{n+1} = N_n + 1,\ N_1 = 1$, as in considered explicit calculation for E_2. In this case, $E_k(N+1) = E_k(N) + (N+1)^k$.

We propose the following compact form for E_k

$$E_k(N) = \frac{d^k}{dx^k} P(x,N)|_{x\Rightarrow 0} \equiv D^k P = P^{(k)}(0,N),$$
$$P(x,N) = \sum_{n=1}^{N} e^{nx} = \frac{e^{(N+1)x} - e^x}{e^x - 1} \tag{144}$$

We take also the following slightly simpler form of $P(x,N)$, for $k = 1, 2, 3, ...$

$$P(x,N) = \sum_{n=0}^{N} e^{nx} = \frac{e^{(N+1)x} - 1}{e^x - 1} \tag{145}$$

As an example, let us calculate $E_1(N)$,

$$E_1(N) = \frac{(N+1)e^{(N+1)x}}{e^x-1} - \frac{(e^{(N+1)x}-1)e^x}{(e^x-1)^2}$$
$$= \frac{(N+1)e^{(N+1)x}(e^x-1) - (e^{(N+1)x}-1)e^x}{(e^x-1)^2} \Downarrow$$
$$\frac{(N+1)(1+(N+1)x...)(x+\frac{x^2}{2}...) - ((N+1)x + \frac{(N+1)^2x^2}{2}...)(1+x...)}{(x+...)^2}$$
$$= (N+1)^2 + (N+1)/2 - (N+1) - (N+1)^2/2 = N(N+1)/2 \qquad (146)$$

We can present the derivative operator in the complex integral form

$$f^{(k)}(0) = \frac{k!}{2\pi i} \oint \frac{dz f(z)}{z^{k+1}} \tag{147}$$

In this form the calculation gives

$$\begin{aligned} S(1,N) &= \frac{1}{2\pi i} \oint \frac{dz}{z^2} \frac{(N+1)z + (N+1)^2 z^2/2}{z + z^2/2} \\ &= \frac{1}{2\pi i} \oint \frac{dz}{z^2} \frac{(N+1) + (N+1)^2 z/2}{1 + z/2} \\ &= \frac{1}{2\pi i} \oint \frac{dz}{z} \frac{(N+1)}{z} (1 - z/2) + (N+1)^2/2 = N(N+1)/2 \end{aligned} \tag{148}$$

By this example we see that the second form of calculation is easier.

Let us consider the integer derivatives of the monomials

$$\begin{aligned}\frac{d^n}{dx^n}x^m &= m(m-1)...(m-(n-1))x^{m-n}, \quad n \leq m, \\ &= \frac{\Gamma(m+1)}{\Gamma(m+1-n)}x^{m-n}. \end{aligned} \tag{149}$$

L.Euler (1707 - 1783) invented the following definition of the fractal derivatives,

$$\frac{d^\alpha}{dx^\alpha}x^\beta = \frac{\Gamma(\beta+1)}{\Gamma(\beta+1-\alpha)}x^{\beta-\alpha}. \tag{150}$$

J.Liouville (1809-1882) takes exponents as a base functions,

$$\frac{d^\alpha}{dx^\alpha}e^{ax} = a^\alpha e^{ax}. \tag{151}$$

The following Cauchy formula

$$I^n_{0,x}f = \int_0^x dx_n \int_0^{x_{n-1}} dx_{n-2} \ldots \int_0^{x_2} dx_1 f(x_1) = \frac{1}{\Gamma(n)} \int_0^x dy (x-y)^{n-1} f(y)$$

permits analytic extension from integer n to complex α,

$$I^\alpha_{0,x}f = \frac{1}{\Gamma(\alpha)} \int_0^x dy (x-y)^{\alpha-1} f(y) \tag{153}$$

J.H. Holmgren invented (in 1863) the following integral transformation,

$$D^{-\alpha}_{c,x} f = \frac{1}{\Gamma(\alpha)} \int_c^x |x-t|^{\alpha-1} f(t) dt. \tag{154}$$

It is easy to show that

$$\begin{aligned} D^{-\alpha}_{c,x} x^m &= \frac{\Gamma(m+1)}{\Gamma(m+1+\alpha)} (x^{m+\alpha} - c^{m+\alpha}), \\ D^{-\alpha}_{c,x} e^{ax} &= a^{-\alpha}(e^{ax} - e^{ac}), \end{aligned} \tag{155}$$

so, $c = 0$, when $m + \alpha \geq 0$, in Holmgren's definition of the fractal calculus, corresponds to the Euler's definition, and $c = -\infty$, when $a > 0$, corresponds to the Liouville's definition.
Holmgren's definition of the fractal calculus reduce to the Euler's definition for finite c, and to the Liouvill's definition for $c = \infty$,

$$\begin{aligned} D^{-\alpha}_{c,x} f &= D^{-\alpha}_{0,x} f - D^{-\alpha}_{0,c} f, \\ D^{-\alpha}_{\infty,x} f &= D^{-\alpha}_{-\infty,x} f - D^{-\alpha}_{-\infty,\infty} f, \\ D^{-\alpha}_{-\infty,x} f &= D^{\alpha}_{0,x} f - D^{-\alpha}_{0,-\infty} f. \end{aligned} \tag{156}$$

We considered the following modification of the $c = 0$ case [Makhaldiani, 2003],

$$D_{0,x}^{-\alpha} f = \frac{|x|^\alpha}{\Gamma(\alpha)} \int_0^1 |1-t|^{\alpha-1} f(xt) dt, = \frac{|x|^\alpha}{\Gamma(\alpha)} B(\alpha, \partial x) f(x)$$
$$= |x|^\alpha \frac{\Gamma(\partial x)}{\Gamma(\alpha + \partial x)} f(x), \ f(xt) = t^{x \frac{d}{dx}} f(x). \quad (157)$$

As an example, consider Euler B-function,

$$B(\alpha, \beta) = \int_0^1 dx |1-x|^{\alpha-1} |x|^{\beta-1} = \Gamma(\alpha)\Gamma(\beta) D_{01}^{-\alpha} D_{0x}^{1-\beta} 1 = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)} \quad (158)$$

Negative Binomial Distribution

Negative binomial distribution (NBD) is defined as

$$P(n) = \frac{\Gamma(n+r)}{n!\Gamma(r)} p^n (1-p)^r, \ \sum_{n\geq 0} P(n) = 1, \tag{159}$$

The Bose-Einstein distribution is a special case of NBD with $r = 1$.
NBD provides a very good parametrization for multiplicity distributions in e^+e^- annihilation; in deep inelastic lepton scattering; in proton-proton collisions; in proton-nucleus scattering. Hadronic collisions at high energies (LHC) lead to charged multiplicity distributions whose shapes are well fitted by a single NBD in fixed intervals of central (pseudo)rapidity η.

A Bose-Einstein, or geometrical, distribution is a thermal distribution for single state systems. An useful property of the negative binomial distribution with parameters

$$< n >,\ k$$

is that it is (also) the distribution of a sum of k independent random variables drawn from a Bose-Einstein distribution with mean $< n > /k$,

$$P_n = \frac{1}{< n > +1}(\frac{< n >}{< n > +1})^n$$
$$= (e^{\beta\hbar\omega/2} - e^{-\beta\hbar\omega/2})e^{-\beta\hbar\omega(n+1/2)},\ T = \frac{\hbar\omega}{\ln\frac{<n>+1}{<n>}}$$
$$\sum_{n\geq 0} P_n = 1,\ \sum nP_n =< n >= \frac{1}{e^{\beta\hbar\omega-1}},\ T \simeq \hbar\omega < n >,\ < n >\gg 1,$$
$$P(x) = \sum_n x^n P_n = (1+ < n > (1-x))^{-1}. \qquad (160)$$

Indeed, for

$$n = n_1 + n_2 + ... + n_k, \tag{161}$$

with n_i independent of each other, the probability distribution of n is

$$P_n = \sum_{n_1,...,n_k} \delta(n - \sum n_i) p_{n_1} ... p_{n_k},$$
$$P(x) = \sum_n x^n P_n = p(x)^k \tag{162}$$

This has a consequence that an incoherent superposition of N emitters that have a negative binomial distribution with parameters $k, < n >$ produces a negative binomial distribution with parameters $Nk, N < n >$.
So, for the GF of NBD we have (N=2)

$$F(k, < n >) F(k, < n >) = F(2k, 2 < n >) \tag{163}$$

And more general formula (N=m) is

$$F(k, < n >)^m = F(mk, m < n >) \tag{164}$$

We can put this equation in the closed nonlocal form

$$Q_q F = F^q, \tag{165}$$

where

$$Q_q = q^D, \quad D = \frac{kd}{dk} + \frac{< n > d}{d < n >} = \frac{x_1 d}{dx_1} + \frac{x_2 d}{dx_2} \tag{166}$$

Note that temperature defined in (160) gives an estimation of the Glukvar temperature when it radiates hadrons. If we take $\hbar\omega = 100MeV$, to $T \simeq T_c \simeq 200MeV$ corresponds $< n > \simeq 1.5$ If we take $\hbar\omega = 10MeV$, to $T \simeq T_c \simeq 200MeV$ corresponds $< n > \simeq 20$. A singular behavior of $< n >$ may indicate corresponding phase transition and temperature. At that point we estimate characteristic quantum $\hbar\omega$.
We see that universality of NBD in hadron-production is similar to the universality of black body radiation.

The Riemann zeta function $\zeta(s)$ is defined for complex $s = \sigma + it$ and $\sigma > 1$ by the expansion

$$\zeta(s) = \sum_{n\geq 1} n^{-s},\ Re\ s > 1,$$
$$= \delta_x^{-s} \frac{x}{1-x}|_{x\rightarrow 1} = \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} e^{-\delta_x t} \frac{x}{1-x}|_{x\rightarrow 1}$$
$$= \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} e^{t\partial_\tau} \frac{1}{e^\tau - 1}|_{\tau\rightarrow 0} = \frac{1}{\Gamma(s)} \int_0^\infty \frac{t^{s-1}dt}{e^t - 1},\ x = e^{-t} \quad (167)$$

All complex zeros, $s = \alpha + i\beta$, of $\zeta(\sigma + it)$ function lie in the critical stripe $0 < \sigma < 1$, symmetrically with respect to the real axe and critical line $\sigma = 1/2$. So it is enough to investigate zeros with $\alpha \leq 1/2$ and $\beta > 0$. These zeros are of three type, with small, intermediate and big ordinates.

The Riemann hypothesis states that the (non-trivial) complex zeros of $\zeta(s)$ lie on the critical line $\sigma = 1/2$.
At the beginning of the XX century Polya and Hilbert made a conjecture that the imaginary part of the Riemann zeros could be the oscillation frequencies of a physical system (ζ - (mem)brane).
After the advent of Quantum Mechanics, the Polya-Hilbert conjecture was formulated as the existence of a self-adjoint operator whose spectrum contains the imaginary part of the Riemann zeros.
The Riemann hypothesis (RH) is a central problem in Pure Mathematics due to its connection with Number theory and other branches of Mathematics and Physics.

The functional equation is

$$\zeta(1-s) = \frac{2\Gamma(s)}{(2\pi)^s}\cos(\frac{\pi s}{2})\zeta(s) \tag{168}$$

From this equation we see the real (trivial) zeros of zeta function:

$$\zeta(-2n) = 0,\ n = 1, 2, ... \tag{169}$$

Also, at s=1, zeta has pole with reside 1.
From Field theory and statistical physics point of view, the functional equation (168) is duality relation, with self dual (or critical) line in the complex plane, at $s = 1/2 + i\beta$,

$$\zeta(\frac{1}{2} - i\beta) = \frac{2\Gamma(s)}{(2\pi)^s}\cos(\frac{\pi s}{2})\zeta(\frac{1}{2} + i\beta), \tag{170}$$

we see that complex zeros lie symmetrically with respect to the real axe.
On the critical line, (nontrivial) zeros of zeta corresponds to the infinite value of the free energy,

$$F = -T\ln\zeta. \tag{171}$$

At the point with $\beta = 14.134725...$ is located the first zero. In the interval $10 < \beta < 100$, zeta has 29 zeros. The first few million zeros have been computed and all lie on the critical line. It has been proved that uncountably many zeros lie on critical line.
The first relation of zeta function with prime numbers is given by the following formula,

$$\zeta(s) = \prod_p (1 - p^{-s})^{-1}, \ Re\ s > 1. \tag{172}$$

Another formula, which can be used on critical line, is

$$\begin{aligned} \zeta(s) &= (1 - 2^{1-s})^{-1} \sum_{n \geq 1} (-1)^{n+1} n^{-s} \\ &= \frac{1}{1 - 2^{1-s}} \frac{1}{\Gamma(s)} \int_0^\infty \frac{t^{s-1} dt}{e^t + 1}, \ Re\ s > 0 \end{aligned} \tag{173}$$

Let us consider the values $q = n, n = 1, 2, 3, ...$ and take sum of the corresponding equations (165), we find

$$\zeta(-D)F = \frac{F}{1-F} \tag{174}$$

In the case of the NBD we know the solutions of this equation.
Now we invent a Hamiltonian H with spectrum corresponding to the set of nontrivial zeros of the zeta function, in correspondence with Riemann hypothesis,

$$-D_n = \frac{n}{2} + iH_n, \;\; H_n = i(\frac{n}{2} + D_n),$$

$$D_n = x_1\partial_1 + x_2\partial_2 + ... + x_n\partial_n, \;\; H_n^+ = H_n = \sum_{m=1}^{n} H_1(x_m),$$

$$H_1 = i(\frac{1}{2} + x\partial_x) = -\frac{1}{2}(x\hat{p} + \hat{p}x), \;\; \hat{p} = -i\partial_x \tag{175}$$

The Hamiltonian $H = H_n$ is hermitian, its spectrum is real. The case $n = 1$ corresponds to the Riemann hypothesis.

The case $n = 2$, corresponds to NBD,

$$\zeta(1+iH_2)F = \frac{F}{1-F},\ \zeta(1+iH_2)|_F = \frac{1}{1-F},$$
$$F(x_1,x_2;h) = (1+\frac{x_1}{x_2}(1-h))^{-x_2} \tag{176}$$

Let us scale $x_2 \to \lambda x_2$ and take $\lambda \to \infty$ in (176), we obtain

$$\zeta(\frac{1}{2}+iH_1(x))e^{-(1-h)x} = \frac{1}{e^{(1-h)x}-1},$$
$$\frac{1}{\zeta(\frac{1}{2}+iH(x))}\frac{1}{e^{\varepsilon x}-1} = e^{-\varepsilon x},$$
$$H(x) = i(\frac{1}{2}+x\partial_x) = -\frac{1}{2}(x\hat{p}+\hat{p}x),\ H^+ = H, \varepsilon = 1-h. \tag{177}$$

Let us take an eigenvector $|n>$ with eigenvalue E_n of H, than

$$< n|\zeta(\frac{1}{2}+iH(x))e^{-(1-h)x} >= \zeta(\frac{1}{2}+iE_n(x)) < n|e^{-(1-h)x} >$$
$$=< n|\frac{1}{e^{(1-h)x}-1} > \tag{178}$$

For zeros of Zeta function, E_n, the eigenfunctions fulfils the following conditions

$$< n|\frac{1}{e^{(1-h)x}-1} >= 0,\ < n|e^{-(1-h)x} >\neq 0. \tag{179}$$

For eigenvalues of H, we have

$$H|n>=E_n|n>,\ H=i(\frac{1}{2}+x\partial_x),\ |n>\sim x^{s_n},\ s_n=-\frac{1}{2}-iE_n,$$
$$<n|\frac{1}{e^{(1-h)x}-1}>\sim\zeta(\frac{1}{2}+iE_n),$$
$$<n|e^{-(1-h)x}>\sim\Gamma(\frac{1}{2}+iE_n). \qquad (180)$$

Brodsky S J, de Tèramond G F, Deur A// Phys. Rev. D 2010. V. 81 P. 096010.

Collins J.C. Renormalization. London: Cambrige Univ. Press, 1984.

C.R.Das, N.Makhaldiani, Work in progress

Diakonov D// Prog. Par. Nucl. Phys. 2003. V. 51 P. 173.

Duff M, Okun L, Veneziano G// J. High Energy Phys. 2002. V. 0203 P. 023.

Ketov S.V. Quantum Non-linear Sigma-models. Springer-Verlag, 2000.

Neal Koblitz, p-adic numbers, p-adic analysis, and Zeta-functions, Springer-Verlag, New York Heidelberg Berlin, 1977.

Maldacena J M // Adv. Theor. Math. Phys. 1988. V. 2 P. 231.

N. Makhaldiani, *Adelic Universe and Cosmological Constant*,
JINR Communications, **E2-2003-215** Dubna 2003, [arXiv:hep-th/0312291].

Makhaldiani N.// Fractal Calculus (H) and some Applications, Physics of Particles and Nuclei Letters. 2011. V. 8. P. 325.

Makhaldiani N Regular method of construction of the reversible dynamical systems and their linear extensions - Quanputers// Atomic Nuclei. 2011. V. 74 P. 1040.

Makhaldiani N Renormdynamics, Unified Field Theories and Universal Distributions of the Multipartical Production Processes// Physics of Particles and Nuclei Letters. 2014. V. 11 P. 1019.

R. Oehme and W. Zimmermann, Commun. Math. Phys. 97 (1985) 569.

L. Okun, The fundamental constants of physics, Sov. Phys. Usp. 34 (1991) 818.

Planck M// Ann. d. Phys. 1900. V. 1 P. 69.

Stoney G. On The Physical Units of Nature, Phil.Mag. 11, 381-391, 1881

Voloshin M.B., Ter-Martyrosian K.A. Gauge Theory of Elementary Particles. Moscow: Atomizdat, 1984.

S. Weinberg, *Gravitation and Cosmology*, New York 1972.

W. Zimmermann, Commun. Math. Phys. 97 (1985) 211

O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri and Y. Oz, Large N field theories, string theory and gravity, *Phys. Rept.* **323** (2000) 183, hep-th/9905111.

K. Aamodt et al. [ALICE collaboration] *Eur. Phys. J.* **C65** (2010) 111 [arXiv:1004.3034], [arXiv:1004.3514].

M. Anselmino, B.L. Ioffe and E. Leader, Sov. J. Nucl. 49, 136, (1989)

H. Aref, Ann. Rev. Fluid Mech. **15** (1983) 345.

V.I. Arnold, *Mathematical Methods of Classical Mechanics*, Springer, New York 1978.

J. Bagger, N. Lambert, *Modeling multiple M2's*, Phys. Rev. D **75** (2007) 045020; [arXiv:hep-th/0611108].

V.V. Balandin, N.I. Golubeva, *Hamiltonian Methods for the Study of Polarized Proton Beam Dynamics in Accelerators and Storage Rings*, [arXiv://arxiv.org/abs/physics/9903032v1].

V. Bargmann, L. Michel, V.L. Telegdi, *Precession of the polarization of particles moving in a homogeneous electromagnetic field*, Phys. Rev. Lett., **2(10)** (1959) 435.

G. Benenti, G. Casati, G. Strini, *Principles of quantum computation and information*, Vol. I: Basic concepts, World Scientific, Singapore 2004; Vol. II: *Basic tools and special topics* World Scientific, Singapore 2007.

J. Beringer et al. (Particle Data Group) *Review of Particle Physics*, Phys. Rev. D **86** (2012) 010001 [1528 pages]

J. D. Bjorken, Phys. Rev. 148, 1467 (1966).

F.A. Berezin, *Introduction to Superanalysis*, Reidel, Dordrecht 1987.

F.A. Berezin, M.S. Marinov, Ann. Phys. (N.Y.) **104** (1977) 336.

N.N. Bogoliubov and D.V. Shirkov, *Introduction to the Theory of Quantized Fields*, New York 1959.

S. Brodsky, G. Farrar, Phys. Rev. Lett. **31** 1153 (1973).

Stanley J. Brodsky, Guy F. de Tèramond, Alexandre Deur, Nonperturbative QCD Coupling and its β function from Light-Front Holography, PRD 81,096010 (2010), arXiv:1002.3948

V. D. Burkert, Phys. Rev. D 63, 097904 (2001).

C. Buttin, C.R. Acad. Sci. Paris. **269** (1969) 87.

A. H. Castro Neto, F. Guinea, N. M. R. Peres, K. S. Novoselov and A. K. Geim, The electronic properties of graphene, Rev. Mod. Phys. 81 , 109-162 (2009).

W. E. Caswell, Phys. Rev. Lett. 33 (1974) 244.

J.C. Collins, *Renormalization*, Cambrige Univ. Press, London 1984.

Ya.Z. Darbaidze, N.V. Makhaldiani, A.N. Sisakian, L.A. Slepchenko, *TMF* **34** (1978) 303.

D. Diakonov, Prog. Par. Nucl. Phys. 51 (2003) 173.

P.A.M. Dirac, *Proc. Roy, Soc.* **A167** (1938) 148.

A.D. Dolgov, Light millicharged particles and large scale cosmic magnetic fields, arXiv:1404.5820v1 [astro-ph.CO] 23 Apr 2014

S. D. Drell, A. C. Hearn, *Exact Sum Rule For Nucleon Magnetic Moments*, *Phys. Rev. Lett.* **16** (1966) 908.

E. Egorian and O. V. Tarasov, Theor. Math. Phys. 41 (1979) 863 [Teor. Mat. Fiz. 41 (1979) 26]

W. Ernst, I. Schmitt, *Nuovo Cim.* **A33** (1976) 195.


L.D. Faddeev, R. Jackiw, Phys.Rev.Lett. **60** (1988) 1692.


L.D. Faddeev and L.A. Takhtajan, *Hamiltonian methods in the theory of solitons*, Springer, Berlin 1990.


I. V. Fialkovsky, D. V. Vassilevich , Quantum Field Theory in Graphene, arXiv:1111.3017


C. S. Fischer and H. Gies, JHEP 0410 (2004) 048.


V. Flambaum, I. Khriplovich and O. Sushkov, Nucl.Phys. A449, p. 750 (1986).


S.B. Gerasimov, *A Sum Rule for Magnetic Moments and Damping of the Nucleon Magnetic Moment in Nuclei*, *J.Nucl.Phys.(USSR)* **2** (1965) 598.


V. L. Ginzburg, *Theoretical Physics and Astrophysics*, Pergamon, New York 1979.


M.L. Goldberger, S.B. Treiman, Phys. Rev. **110** 1178 (1958).


D.J. Gross, F. Wilczek, Phys. Rev. Lett. **30** 1343 (1973).


S. S. Gubser, I. R. Klebanov and A. M. Polyakov, Phys. Lett. B 428, 105 (1998) [arXiv:hep-th/9802109];


A. Gustavsson, *Algebraic structures on parallel M2-branes*,
Nucl. Phys. B.**811** (2009) 66; [arXiv:0709.1260 [hep-th]].


G. Gutin, A.P. Punnen (Eds), *The Traveling Salesman Problem and Its Variations*, Combinatorial Optimization Series, Kluwer, Boston 2002.


W. Heisenberg, *Introduction to the Unified field Theory of Elementary particles,* Interscience Publishers, London 1966.


G. 't Hooft, report at the *Marseille Conference on Yang-Mills Fields*, 1972.


G. 't Hooft, *Nucl.Phys.* **B 61** (1973) 455.


Ernst-Michael Ilgenfritz, Axel Maas, Topological aspects of G2 Yang-Mills theory, arXiv:1210.5963.


J.D. Jackson, *Classical Electrodynamics, 3rd ed.* JohnWiley & Sons, Inc. New York 1999.


X. Ji, J.Osborne, J.Phys. G27 127 (2001).


D. R. T. Jones, Nucl. Phys. B 75 (1974) 531;


M. Kaku, *Strings, Conformal Fields, and M-Theory*, Springer, New York 2000.


A. L. Kataev, S. A. Larin, Analytical five-loop expressions for the renormalization group QED β-function in different renormalization schemes, Pisma v ZhETF, vol.96, iss. 1, (2012), pp.64-67


D.I. Kazakov, *Supersymmetric Generalization of the Standard Model of Fundamental Interactions*, Textbook, JINR Dubna 2004.


D.I. Kazakov, D.V. Shirkov, Fortschr. d. Phys. **28** 447 (1980).


D.I. Kazakov, L.R. Lomidze, N.V. Makhaldiani, A.A. Vladimirov, *Ultraviolet Asymptotics in Renormalizable Scalar Theories*, JINR Communications, **E2-8085**, Dubna 1974.


A.Yu. Kitaev, A. Shen, M.N. Vyalyi, *Classical and Quantum Computation*, American Mathematical Society, 2002.


Z. Koba, H.B. Nielsen, P. Olesen, *Nucl. Phys.* **B40** (1972) 317.


N.V. Makhaldiani, *Approximate methods of the field theory and their applications in physics of high energy, condensed matter, plasma and hydrodynamics*, Dubna 1980.


S. A. Larin and J. A. M. Vermaseren, Phys. Lett. B 303 (1993) 334.


C. Lerche and L. von Smekal, Phys. Rev. D 65 (2002) 125006.


N.M. Makhaldiani, *Computational Quantum Field Theory*, JINR Communication, **P2-86-849**, Dubna 1986.


N.V. Makhaldiani, *A New Approach to the Problem of Space Compactification,*
JINR Communications, **P2-87-306**, Dubna 1987.


N.V. Makhaldiani, *Number Fields Dynamics and the Compactification of Space Problem in the Unified Theories of Fields and Strings*, JINR Communications, **P2-88-916**, Dubna 1988.


N. Makhaldiani, *The System of Three Vortexes of Two-Dimensional Ideal Hydrodynamics as a New Example of the (Integrable) Nambu-Poisson Mechanics*, JINR Communications **E2-97-407**, Dubna 1997; [arXiv:solv-int/9804002].


N. Makhaldiani, *The Algebras of the Integrals of Motion and Modified Bochner-Killing-Yano Structures of the Point particle Dynamics*, JINR Communications **E2-99-337**, Dubna 1999.


N. Makhaldiani, *New Hamiltonization of the Schrödinger Equation by Corresponding Nonlinear Equation for the Potential*, JINR Communications, **E2-2000-179**, Dubna 2000.


N. Makhaldiani, *How to Solve the Classical Problems on Quantum Computers*, JINR Communications, **E2-2001-137**, Dubna 2001.


N. Makhaldiani, *Classical and Quantum Problems for Quanputers*, [arXiv:quant-ph/0210184].


N. Makhaldiani, *Adelic Universe and Cosmological Constant*, JINR Communications, **E2-2003-215** Dubna 2003, [arXiv:hep-th/0312291].


N. Makhaldiani, *Nambu-Poisson dynamics of superintegrable systems*, *Atomic Nuclei*, **70** (2007) 564.


N. Makhaldiani, *Theory of Quanputers*, Sovremennaia Matematica i ee Prilozhenia, **44** (2007) 113; Journal of Mathematical Sciences, **153** (2008) 159.


N.V. Makhaldiani, *Renormdynamics and Scaling Functions*, in Proc. of the *XIX International Baldin Seminar on High Energy Physics Problems* eds. A.N.Sissakian, V.V.Burov, A.I.Malakhov, S.G.Bondartenko, E.B.Plekhanov, Vol.II, p. 175, Dubna 2008.


N.V. Makhaldiani, *Renormdynamics, multiparticle production, negative binomial distribution and Riemann zeta function*, [arXiv:1012.5939v1 [math-ph]] 24 Dec 2010.


N. Makhaldiani, *Fractal Calculus (H) and some Applications*, *Physics of Particles and Nuclei Letters*, **8** 325 (2011).


N. Makhaldiani, *Regular method of construction of the reversible dynamical systems and their linear extensions - Quanputers*, Atomic Nuclei, **74** (2011) 1040.


Nugzar Makhaldiani, *Nambu-Poisson Dynamics with Some Applications*,
Physics of Particles and Nuclei, **43** (2012) 703.


Nugzar Makhaldiani, *Renormdynamics, coupling constant unification and universal properties of the multiparticle production*, XXI International Baldin Seminar on High Energy Physics Problems, September 10-15, 2012, PoS(Baldin ISHEPP XXI)068.


N. V. Makhaldiani, *Renormdynamics, Multiparticle Production, Negative Binomial Distribution, and Riemann Zeta Function*, Physics of Atomic Nuclei, **76** 1169 (2013).


N. Makhaldiani, O. Voskresenskaya, *On the correspondence between the dynamics with odd and even brackets and generalized Numbu's mechanics,*
JINR Communications, **E2-97-418**, Dubna 1997.


J. M. Maldacena, Adv. Theor. Math. Phys. 2, 231 (1998) [Int. J. Theor. Phys. 38, 1113 (1999)] [arXiv:hep-th/9711200].


V. Matveev, R. Muradyan, A. Tavkhelidze, Lett. Nuovo Cimento **7** 719 (1973).


V.A. Matveev, A.N. Sisakian, L.A. Slepchenko, *Nucl. Phys.* **23** (1976) 432.


A.V. Meleshko, N.N. Konstantinov, *Dynamics of vortex systems*,
Naukova Dumka, Kiev 1993.


Y. Nambu, Phys.Rev. D **7** (1973) 2405.


M.A. Nielsen, I.L. Chuang, *Quantum computation and quantum information*, Cambridge University Press, Cambridge 2000.


J. M. Pawlowski, D. F. Litim, S. Nedelko and L. von Smekal, Phys. Rev. Lett. 93 (2004) 152002.


H.D. Politzer, Phys. Rev. Lett. **30** 1346 (1973).


M.C.M. Rentmeester, R.G.E. Timmermans, J.L. Friar, J.J. de Swart, Phys. Rev. Lett. **82** 4992 (1999).

T. van Ritbergen, J. A. M. Vermaseren and S. A. Larin, Phys. Lett. B 400 (1997)379.

B. L. Roberts and W. J. Marciano (eds.), Lepton Dipole Moments, Advanced Series on Directions in High Energy Physics, Vol. 20 (World Scientific, 2010).

A. Sakharov, Pisma Zh.Eksp.Teor.Fiz. 5, 32 (1967), Reprinted in Sov. Phys. Usp. 34 (1991) 392-393 [Usp. Fiz. Nauk 161 (1991) No. 5 61-64].

A. Samarskii, A. Gulin, *Numerical Methods*, Nauka, Moscow 1989.

J. Schwinger, *Magnetic Charge and Quantum Field Theory, Phys. Rev.* **144** (1966) 1087.

Ya.G. Sinai, *Topics in Ergodic Theory*, Princeton University Press, Princeton NJ 1993.

O. V. Tarasov, A. A. Vladimirov and A. Y. Zharkov, Phys. Lett. B 93 (1980) 429.

Norman Margolus, Tommaso Toffoli, *Cellular Automaton Machines*,
MIT Press, Cambridge 1987.

M.V. Tokarev, I. Zborovský, *Z-Scaling in the Proton-Proton Collisions at RHIC*, in *Investigations of Properties of Nuclear Matter at High Temperature and Densities, Editid by A.N. Sisakian, F.A. Soifer*, Dubna 2007

L.H. Thomas, Philos. Mag. **3** (1927) 1.

M.B. Voloshin, K.A. Ter-Martyrosian, *Gauge Theory of Elementary Particles*, Atomizdat, Moscow 1984.

S. Weinberg, AIP Conf.Proc. 272, 346 (1993).

E.T. Whittaker, *A Treatise on the Analytical Dynamics*, Cambridge 1927.

E. Witten, Adv. Theor. Math. Phys. 2, 253 (1998) [arXiv:hep-th/9802150].

Zelikin, Mikhail I. (2000), Homogeneous Spaces and the Riccati Equation in the Calculus of Variations, Berlin: Springer-Verlag

D. Zwanziger, Phys. Rev. D 65 (2002) 094039.